

11/23/2015; Page 1

٦

# SUGGESTED FORMULATION

Ingredient Listing	Qty.	Unit	NDC #	Supplier	Lot Number	Expiry Date
APIs**	TBD					
Glycerin, USP***	6.0	mL				
Tuna Flavor (Powder)	1.50	g				
Chew-A-Treat <sup>™</sup> Compound A	23.10	g				
Chew-A-Treat <sup>™</sup> Compound C	46.21	g				

\*\*Note: APIs can be added by using 0.7 displacement factor to accordingly reduce Compound A and C in the 1:2 ratio. \*\*\*Note: Caprylic/Capric Triglyceride can be used to replace Glycerin to reduce stickiness.

# SPECIAL PREPARATORY CONSIDERATIONS

Ingredient-Specific Information	
Hygroscopic (protect from moi	sture whenever possible): Glycerin
Suggested Preparatory Guidelines	
Non-Sterile Preparat	on Sterile Preparation
<u>Processing Error /</u> <u>Testing Considerations</u> :	To account for processing error considerations during preparation, it is suggested to measure an additional <b>5 to 9%</b> of the required quantities of ingredients.
Special Instruction:	Protective apparel, such as a lab coat, disposable gloves, eyewear and face-masks should always be worn.
	This procedure requires the use of very small quantities of ingredients. All calculations and preparation techniques must be verified before dispensing the final product.



11/23/2015; Page 2

Suggested Formula	Blank Oral Chewable Treats (Solid Suspension, 60 x 1.0 mL Treats)	FIN	F 006 536	
----------------------	---	-----	-----------	--

# SUGGESTED PREPARATION (for 60 x 1.0 mL Treats)

Weigh and / or measure the following ingredients when appropriate:

Ingredient Listing	Qty.	Unit	Multiplication factor <sup>(*)</sup> :	Processing Error	Qty. to measure
APIs	TBD				
Glycerin, USP §	6.0	mL			
Tuna Flavor (Powder)	1.50	g			
Chew-A-Treat <sup>™</sup> Compound A	23.10	g			
Chew-A-Treat <sup>™</sup> Compound C	46.21	g	8		

\* Takes into account increased batch size conversions and density conversions, if required.

§ Weigh / measure just prior to use.

### **Preparatory Instruction**

## 1. Mold lubrication:

A. Lubricate all parts of the Chew-A-Treat<sup>™</sup> Mold with suitable vegetable spray and set aside.

<u>Note</u>: Selected vegetable spray needs to be compatible with API(s) and all other ingredients within the formulation.

## 2. **Powder preparation:**

- A. Triturate the Chew-A-Treat<sup>™</sup> Compound C to form a fine, homogeneous powder blend.
- B. Triturate the APIs to form a fine, homogeneous powder blend.
- C. By geometric addition, combine and mix the following ingredients together to form a homogeneous mixture:

-Fine, homogeneous powder blend (Step 2B)-Tuna Flavor (Powder)-Fine, homogeneous powder blend (Step 2A)

D. Transfer the mixture (Step 2C) to a beaker and incrementally add the Glycerin.

Specifications: Continuously mix until homogeneous.

End result: Homogeneous moist blend.

<u>Note</u>: The mixture will have a sand-like texture. If it begins to stick to your stirring device or beaker, you can simply scrape it off and continue mixing.



11/23/2015; Page 3

Suggested Formula		Blank Oral Chewable Treats (Solid Suspension, 60 x 1.0 mL Treats)	FIN	F 006 536	
3.	Medium integration:				
	A. P	A. Prepare a hot water bath to between 60°C and 65°C.			
	B. U	Jsing the hot water bath, melt the Chew-A-Treat <sup>™</sup> Compound A.			
	<u>S</u>	pecifications: Continuously mix. Maintain temperature between 60°C and 65°C.			
	Ē	and result: Homogeneous thick, sticky liquid-like dispersion.			
		ncrementally add the homogeneous moist blend (Step 2D) to the melted Chew-A-Treat <sup>Th</sup> Step 3B).	<sup>м</sup> Com	pound A	
	<u>S</u>	pecifications: Continuously mix until homogeneous. Maintain temperature between 60°C and 65°C.			
	Ē	and result: Homogeneous thick, sticky paste-like dispersion.			
4.	Mold filling:				
	A. P	properly lubricate gloves (thumbs, forefingers, and palms) throughout filling.			
	r	B. Continuing to heat the mixture between 60 to 65°C, remove a marble-sized quantity of the mixture at a time and roll it between your thumb and forefinger to form a cylinder. Pack the 60 mold cavities with these cylinders and scrape off any excess with pre-lubricated plastic scraper.			
	C. Once the treats have been packed down with an extractor, apply a thin layer of lubricant to give a shiny surface.			a shiny surface.	
		Allow the treats to cool at room temperature for at least 15 minutes and then remove then xtractor.	n from	the mold using the	
5.	Valic	lation technique:			
	A. V	Veigh 12 chewable treats.			
		The final weight of each chewable treat shall not be less than 90% and not more than alculated weight for each unit $(1.31 \text{ g} + (0.3 \times \text{APIs per treat in g})$ in accordance to USP			
6.	Prod	uct transfer:			
	Trans	sfer the final product into the specified dispensing container (see "Packaging Requirement	nts").		



11/23/2015; Page 4

Suggested	Blank Oral Chewable Treats (Solid Suspension, 60 x 1.0 mL Treats)	FIN	F 006 536	1
Formula	Blain of a chevalle freuds (Sona Subpension, of A fit hill freuds)		1 000 000	

SUIGGESTED	PRESENTATION
JUUULUILU	INCOLNIATION

Estima Beyond-Use D		30 days, as per USP.	Packaging Requirements		Manually put into Chew-A-Treat blisters and cold seal with foil labels.
	1	Use as directed. Do not exceed dose.	d prescribed	4	Keep in a dry place.
Auxiliary Labels	2	Keep out of reach of children.		5	Consult your health care practitioner if any prescription or over-the-counter medications are currently being used or are prescribed for future use.
	3 For Feline use only.			6	Keep at room temperature $(20^{\circ}C - 23^{\circ}C)$ .
Pharmacist Instructions	Add any auxiliary labels specific to the API to the dispensing container as deemed necessary.				
Patient Instructions	If allergic reactions occur, consult your pharmacist.				

### REFERENCES

1.	Lozenges/Troches. In: Allen, LV, Jr. <i>The Art, Science and Technology of Pharmaceutical Compounding Third Edition</i> . American Pharmaceutical Association; 2008: 153.
2.	Glycerin. In: Rowe RC. <i>Handbook of Pharmaceutical Excipients</i> , 7 <sup>th</sup> Edition. American Pharmaceutical Association; 2012: 324.
3.	USP <795>. United States Pharmacopeia XXXVIII / National Formulary 33. Rockville, MD. US Pharmacopeial Convention, Inc. 2015: 559.

DISCLAIMER: MEDISCA NETWORK INC., HEREBY REFERRED TO AS 'THE NETWORK', HAS PROVIDED THE FORMULA AND INSTRUCTIONS ABOVE AS A MODEL FOR EDUCATIONAL PURPOSES ONLY ON THE BASIS OF THE RECOGNIZED COMPENDIA AND TEXTS REFERENCED AT THE END OF THIS DOCUMENT. THE NETWORK TAKES NO RESPONSIBILITY FOR THE VALIDITY OR ACCURACY OF THIS INFORMATION OR FOR ITS SAFETY OR EFFECTIVENESS, NOR FOR ANY USE THEREOF, WHICH IS AT THE SOLE RISK OF THE LICENSED PHARMACIST. ADJUSTMENTS MAY BE NEEDED TO MEET SPECIFIC PATIENT NEEDS AND IN ACCORDANCE WITH A LICENSED PHARMACIST. ADJUSTMENTS MAY BE NEEDED TO MEET SPECIFIC PATIENT NEEDS AND IN ACCORDANCE WITH A LICENSED PRESCRIBER'S PRESCRIPTION. THE PHARMACIST MUST EMPLOY APPROPRIATE TESTS TO DETERMINE THE STABILITY OF THIS SUGGESTED FORMULA. THE NETWORK CANNOT BE HELD LIABLE TO ANY PERSON OR ENTITY CONCERNING CLAIMS, LOSS, OR DAMAGE CAUSED BY, OR ALLEGED TO BE CAUSED BY, DIRECTLY OR INDIRECTLY, THE USE OR MISUSE OF THE INFORMATION CONTAINED IN THIS SUGGESTED FORMULA. IN ALL CASES IT IS THE RESPONSIBILITY OF THE LICENSED PHARMACIST TO KNOW THE LAW, TO COMPOUND ANY FINISHED PRODUCT AND TO DISPENSE THESE PRODUCTS IN ACCORDANCE WITH FEDERAL AND STATE LAW.