



MEDISCA® NETWORK INC.  
TECHNICAL SUPPORT SERVICES  
FORMULATION CHEMISTRY DEPARTMENT  
TOLL-FREE: 866-333-7811  
TELEPHONE: 514-905-5096  
FAX: 514-905-5097  
[technicalservices@medisca.net](mailto:technicalservices@medisca.net)

6/7/2018; Page 1

Suggested Formula	Sulfamethoxazole 40 mg/mL, Trimethoprim 8 mg/mL Oral Liquid (Suspension, 150 mL)	FIN	F 007 892
-------------------	--	-----	-----------

### SUGGESTED FORMULATION

Ingredient Listing	Qty.	Unit	NDC #	Supplier	Lot Number	Expiry Date
Sulfamethoxazole, USP	6.000	g				
Trimethoprim, USP	1.200	g				
Medisca Oral Mix (Flavored Suspending Vehicle)	8.0	mL				
Medisca Oral Mix (Flavored Suspending Vehicle)	120.0	mL				
Medisca Oral Mix (Flavored Suspending Vehicle)	q.s. to 150.0	mL				





Suggested Formula	Sulfamethoxazole 40 mg/mL, Trimethoprim 8 mg/mL Oral Liquid (Suspension, 150 mL)	FIN	F 007 892
-------------------	--	-----	-----------

## SPECIAL PREPARATORY CONSIDERATIONS

### Ingredient-Specific Information

**Light Sensitive** (protect from light whenever possible):

*Sulfamethoxazole, Trimethoprim*

### Suggested Preparatory Guidelines

Non-Sterile Preparation     Sterile Preparation

Processing Error / Testing Considerations: To account for processing error considerations during preparation, it is suggested to measure an additional **5 to 9%** of the required quantities of ingredients.

Special Instruction: This formula may contain one or more Active Pharmaceutical Ingredients (APIs) that may be classified as hazardous, please refer & verify the current NIOSH list of Antineoplastic and Other Hazardous Drugs in Healthcare Settings, 2016. **General Chapter <800> Hazardous Drugs – Handling in Healthcare Settings** was formally published February 1, 2016 in the First Supplement to USP 39-NF 34 and has a delayed **official implementation date of December 31<sup>st</sup>, 2019**.

This formula must be prepared within the appropriate facilities under adequate environmental conditions, following the necessary guidelines and procedures as stated within *USP 795* and *USP 800*, when handling hazardous drugs. Only trained and qualified personnel must prepare this formula.

All required personal protective equipment (hazardous if applicable), such as but not limited to, lab coat, protective sleeves, gloves both inner and outer if applicable, dedicated shoe covers, hairnet, beard cover, eyewear, appropriate face mask, respirator and face shield, etc., where applicable must be worn at all times.

If applicable, follow all required procedures for hazardous drug handling including but not limited to procurement, transport, storage, preparation, dispensing, administration, clean up (spills) & disposal.

If you are a registered 503B facility, please refer to all relevant guidance documents including but not limited to the Code of Federal Regulations (CFR), Guidance for Industry (GFIs) and Compliance Policy Guides (CPGs).

This procedure requires the use of very small quantities of ingredients. All calculations and preparation techniques must be verified before dispensing the final product.



Suggested Formula	Sulfamethoxazole 40 mg/mL, Trimethoprim 8 mg/mL Oral Liquid (Suspension, 150 mL)	FIN	F 007 892
-------------------	--	-----	-----------

**SUGGESTED PREPARATION (for 150 mL)**

Weigh and / or measure the following ingredients when appropriate:

Ingredient Listing	Qty.	Unit	Multiplication factor (*): ____	Processing Error	Qty. to measure
Sulfamethoxazole, USP §	6.000	g			
Trimethoprim, USP §	1.200	g			
Medisca Oral Mix (Flavored Suspending Vehicle)	8.0	mL			
Medisca Oral Mix (Flavored Suspending Vehicle)	120.0	mL			
Medisca Oral Mix (Flavored Suspending Vehicle)	q.s. to 150.0	mL			

\* Takes into account increased batch size conversions and density conversions, if required.

§ Weigh / measure just prior to use.

**Preparatory Instruction**

1.	<p><b><u>Powder-liquid preparation:</u></b></p> <p>A. Combine and triturate the following ingredients together to form a fine, homogeneous powder blend:</p> <ul style="list-style-type: none"> <li>-Sulfamethoxazole</li> <li>-Trimethoprim</li> </ul> <p>B. Levigate the fine, homogeneous powder blend (Step 1A) with the Oral Mix (Flavored Suspending Vehicle) (8.0 mL <i>plus</i> processing error adjustments).</p> <p><u>End result:</u> Homogeneous paste-like dispersion.</p>
2.	<p><b><u>Powder-liquid integration:</u></b></p> <p>A. Incrementally add the homogeneous paste-like dispersion (Step 1B) to the Oral Mix (Flavored Suspending Vehicle) (120.0 mL <i>plus</i> processing error adjustments).</p> <p><u>Specifications:</u> Continuously mix.</p> <p><u>End result:</u> Homogeneous liquid-like dispersion.</p>
3.	<p><b><u>Filling to volume:</u></b></p> <p>A. Add additional Oral Mix (Flavored Suspending Vehicle) to the mixture (Step 2A) to fill to the required batch size (150.0 mL <i>plus</i> processing error adjustments).</p> <p><u>Specifications:</u> Continuously mix.</p> <p><u>End result:</u> Homogeneous liquid-like dispersion.</p>



Suggested Formula	Sulfamethoxazole 40 mg/mL, Trimethoprim 8 mg/mL Oral Liquid (Suspension, 150 mL)	FIN	F 007 892
-------------------	--	-----	-----------

4.	<p><b><u>Product transfer:</u></b></p> <p>Transfer the final product into the specified dispensing container (see “Packaging Requirements”).</p> <p><u>Note:</u> Continuously mix the final product during the transfer process in order to maintain homogeneity.</p>
----	---

**SUGGESTED PRESENTATION**

Estimated Beyond-Use Date & Packaging Requirements	Amber PET bottle or PreciseDose Dispenser™ Amber Syringe: 90 days at 5°C or 25°C, based on available stability studies through Medisca*.			
	*Suggested BUD is based on the <b>exact</b> execution of the indicated ingredient list, quantities and procedures listed within this formulation.			
<p><b>Note:</b> <i>This data is provided for informational purposes only, representing the results of a study of the product stability with various active pharmaceutical ingredients. It does not serve, and may not be construed, as a representation or guarantee of product performance. In all cases the practitioner is advised to consult recognized pharmaceutical compendia and other recognized sources for product formulation and other product characteristics, including stability. MEDISCA Network Inc. makes no warranties or representations with regard to the functioning or appropriateness of this product in any compounded formulation, which use is solely at the discretion and liability of the practitioner.</i></p>				
Auxiliary Labels	1	Use as directed. Do not exceed prescribed dose.	6	Cap tightly after use.
	2	Keep out of reach of children.	7	Protect from light.
	3	Do not take with alcohol, sleep aids, tranquilizers or other CNS depressants.	8	Consult your health care practitioner if any prescription or over-the-counter medications are currently being used or are prescribed for future use.
	4	Keep refrigerated (Do not freeze) OR keep at room temperature (25°C).	9	<b>Shake well before use.</b>
	5	Keep in a dry place.	10	
Pharmacist Instructions	Add any auxiliary labels specific to the API to the dispensing container as deemed necessary.			
Patient Instructions	Contact your pharmacist in the event of adverse reactions.			



Suggested Formula	Sulfamethoxazole 40 mg/mL, Trimethoprim 8 mg/mL Oral Liquid (Suspension, 150 mL)	FIN	F 007 892
-------------------	--	-----	-----------

## REFERENCES

1.	Suspensions. In: Allen, LV, Jr. <i>The Art, Science, and Technology of Pharmaceutical Compounding Fifth Edition</i> . American Pharmacists Association; 2016: 279.
2.	Sulfamethoxazole. In: Sweetman SC, ed. <i>Martindale: The Complete Drug Reference, 36<sup>th</sup> Edition</i> . London, England: The Pharmaceutical Press; 2009: 340.
3.	Trimethoprim. In: Sweetman SC, ed. <i>Martindale: The Complete Drug Reference, 36<sup>th</sup> Edition</i> . London, England: The Pharmaceutical Press; 2009: 355.
4.	Sulfamethoxazole (Monograph). In: O'Neil MJ. <i>The Merck Index 15<sup>th</sup> Edition</i> . Whitehouse Station, NJ: Merck & Co, Inc.; 2013: Monograph #9050.
5.	Trimethoprim (Monograph). In: O'Neil MJ. <i>The Merck Index 15<sup>th</sup> Edition</i> . Whitehouse Station, NJ: Merck & Co, Inc.; 2013: Monograph #9878.
6.	Trimethoprim. In: Trissel LA. <i>Trissel's Stability of Compounded Formulations, 5th Edition</i> . American Pharmaceutical Association; 2012: 488.
7.	Sulfamethoxazole (Monograph). <i>United States Pharmacopeia XL / National Formulary 35</i> . Rockville, MD. US Pharmacopeial Convention, Inc. 2017: 6267.
8.	Trimethoprim (Monograph). <i>United States Pharmacopeia XL / National Formulary 35</i> . Rockville, MD. US Pharmacopeial Convention, Inc. 2017: 6597.
9.	USP <795>. <i>United States Pharmacopeia XLI / National Formulary 36</i> . Rockville, MD. US Pharmacopeial Convention, Inc. 2018: 6546.

DISCLAIMER: MEDISCA NETWORK INC., HEREBY REFERRED TO AS 'THE NETWORK', HAS PROVIDED THE FORMULA AND INSTRUCTIONS ABOVE AS A MODEL FOR EDUCATIONAL PURPOSES ONLY ON THE BASIS OF THE RECOGNIZED COMPENDIA AND TEXTS REFERENCED AT THE END OF THIS DOCUMENT. THE NETWORK TAKES NO RESPONSIBILITY FOR THE VALIDITY, SCHEDULING OR ACCURACY OF THIS INFORMATION OR FOR ITS SAFETY OR EFFECTIVENESS, NOR FOR ANY USE THEREOF, WHICH IS AT THE SOLE RISK OF THE LICENSED PHARMACIST. ADJUSTMENTS MAY BE NEEDED TO MEET SPECIFIC PATIENT NEEDS AND IN ACCORDANCE WITH A LICENSED PRESCRIBER'S PRESCRIPTION. THE PHARMACIST MUST EMPLOY APPROPRIATE TESTS TO DETERMINE THE STABILITY OF THIS SUGGESTED FORMULA. THE NETWORK CANNOT BE HELD LIABLE TO ANY PERSON OR ENTITY CONCERNING CLAIMS, LOSS, OR DAMAGE CAUSED BY, OR ALLEGED TO BE CAUSED BY, DIRECTLY OR INDIRECTLY, THE USE OR MISUSE OF THE INFORMATION CONTAINED IN THIS SUGGESTED FORMULA. IN ALL CASES IT IS THE RESPONSIBILITY OF THE LICENSED PHARMACIST TO KNOW THE LAW, TO COMPOUND ANY FINISHED PRODUCT AND TO DISPENSE THESE PRODUCTS IN ACCORDANCE WITH FEDERAL AND STATE LAW.