



MEDISCA® NETWORK INC.
TECHNICAL SUPPORT SERVICES
FORMULATION CHEMISTRY DEPARTMENT
TOLL-FREE: 866-333-7811
TELEPHONE: 514-905-5096
FAX: 514-905-5097
technicalservices@medisca.net

7/26/2018; Page 1

Suggested Formula	Pyridoxine Hydrochloride 20 mg/mL Oral Liquid (Solution, 100 mL)	FIN	F 007 860
-------------------	--	-----	-----------

SUGGESTED FORMULATION

Ingredient Listing	Qty.	Unit	NDC #	Supplier	Lot Number	Expiry Date
Pyridoxine Hydrochloride, USP	2.000	g				
Stevia Powder	0.50	g				
Methylcellulose Gel (1%)	5.0	mL				
Methylcellulose Gel (1%)	q.s. to 100.0	mL				





Suggested Formula	Pyridoxine Hydrochloride 20 mg/mL Oral Liquid (Solution, 100 mL)	FIN	F 007 860
-------------------	--	-----	-----------

SPECIAL PREPARATORY CONSIDERATIONS

Ingredient-Specific Information

Hygroscopic (protect from moisture whenever possible): *Stevia Powder*

Light Sensitive (protect from light whenever possible): *Pyridoxine Hydrochloride*

Suggested Preparatory Guidelines

Non-Sterile Preparation Sterile Preparation

Processing Error / Testing Considerations: To account for processing error considerations during preparation, it is suggested to measure an additional **5 to 9%** of the required quantities of ingredients.

Special Instruction: This formula may contain one or more Active Pharmaceutical Ingredients (APIs) that may be classified as hazardous, please refer & verify the current NIOSH list of Antineoplastic and Other Hazardous Drugs in Healthcare Settings, 2016. **General Chapter <800> Hazardous Drugs – Handling in Healthcare Settings** was formally published February 1, 2016 in the First Supplement to USP 39-NF 34 and has a delayed **official implementation date of December 31st, 2019**.

This formula must be prepared within the appropriate facilities under adequate environmental conditions, following the necessary guidelines and procedures as stated within *USP 795* and *USP 800*, when handling hazardous drugs. Only trained and qualified personnel must prepare this formula.

All required personal protective equipment (hazardous if applicable), such as but not limited to, lab coat, protective sleeves, gloves both inner and outer if applicable, dedicated shoe covers, hairnet, beard cover, eyewear, appropriate face mask, respirator and face shield, etc., where applicable must be worn at all times.

If applicable, follow all required procedures for hazardous drug handling including but not limited to procurement, transport, storage, preparation, dispensing, administration, clean up (spills) & disposal.

If you are a registered 503B facility, please refer to all relevant guidance documents including but not limited to the Code of Federal Regulations (CFR), Guidance for Industry (GFI) and Compliance Policy Guides (CPGs).

This procedure requires the use of very small quantities of ingredients. All calculations and preparation techniques must be verified before dispensing the final product.



Suggested Formula	Pyridoxine Hydrochloride 20 mg/mL Oral Liquid (Solution, 100 mL)	FIN	F 007 860
-------------------	--	-----	-----------

SUGGESTED PREPARATION (for 100 mL)

Weigh and / or measure the following ingredients when appropriate:

Ingredient Listing	Qty.	Unit	Multiplication factor (*): ____	Processing Error	Qty. to measure
Pyridoxine Hydrochloride, USP	2.000	g			
Stevia Powder §	0.50	g			
Methylcellulose Gel (1%)	5.0	mL			
Methylcellulose Gel (1%)	q.s. to 100.0	mL			

* Takes into account increased batch size conversions and density conversions, if required.

§ Weigh / measure just prior to use.

Preparatory Instruction

1.	<p><u>Powder-liquid preparation:</u></p> <p>A. Combine and triturate the following ingredients together to form a fine, homogeneous powder blend:</p> <ul style="list-style-type: none">-Pyridoxine Hydrochloride-Stevia Powder <p>B. Levigate the fine, homogeneous powder blend (Step 1A) with the Methylcellulose Gel (1%) (5.0 mL <i>plus</i> processing error adjustments).</p> <p><u>End result:</u> Homogeneous liquid-like dispersion.</p>
2.	<p><u>Filling to volume:</u></p> <p>A. Add additional Methylcellulose Gel (1%) to the Homogeneous liquid-like dispersion (Step 1B) to fill to the required batch size (100.0 mL <i>plus</i> processing error adjustments).</p> <p><u>Specifications:</u> Continuously mix until homogeneous.</p> <p><u>End result:</u> Homogeneous liquid-like solution.</p>
3.	<p><u>Product transfer:</u></p> <p>A. Transfer the final product into the specified dispensing container (see “Packaging Requirements”).</p>



Suggested Formula	Pyridoxine Hydrochloride 20 mg/mL Oral Liquid (Solution, 100 mL)	FIN	F 007 860
-------------------	--	-----	-----------

SUGGESTED PRESENTATION

Estimated Beyond-Use Date	14 days, refrigerated, as per USP.	Packaging Requirements	- Tightly closed, light resistant dispensing bottle. - To be administered with a metered dose-measuring device.	
Auxiliary Labels	1	Use as directed. Do not exceed prescribed dose.	5	Keep out of reach of children.
	2	Protect from light.	6	Consult your health care practitioner if any prescription or over-the-counter medications are currently being used or are prescribed for future use.
	3	Keep refrigerated. Do not freeze.	7	Cap tightly after use.
	4	Do not take with alcohol, sleep aids, tranquilizers or other CNS depressants.	8	Keep in a dry place.
Pharmacist Instructions	Add any auxiliary labels specific to the API to the dispensing container as deemed necessary.			
Patient Instructions	Contact your pharmacist in the event of adverse reactions.			



Suggested Formula	Pyridoxine Hydrochloride 20 mg/mL Oral Liquid (Solution, 100 mL)	FIN	F 007 860
-------------------	--	-----	-----------

REFERENCES

1.	Solutions. In: Allen, LV, Jr. <i>The Art, Science, and Technology of Pharmaceutical Compounding Fifth Edition</i> . American Pharmacists Association; 2016: 263.
2.	Methylcellulose. In: Rowe RC. <i>Handbook of Pharmaceutical Excipients, 7th Edition</i> . American Pharmaceutical Association; 2012: 496.
3.	Pyridoxine Hydrochloride. In: Sweetman SC, ed. <i>Martindale: The Complete Drug Reference, 36th Edition</i> . London, England: The Pharmaceutical Press; 2009: 1978.
4.	Pyridoxine Hydrochloride (Monograph). In: O'Neil MJ. <i>The Merck Index 15th Edition</i> . Whitehouse Station, NJ: Merck & Co, Inc.; 2013: Monograph #8095.
5.	Pyridoxine Hydrochloride. In: Trissel LA. <i>Trissel's Stability of Compounded Formulations, 5th Edition</i> . American Pharmaceutical Association; 2012: 418.
6.	Pyridoxine Hydrochloride (Monograph). <i>United States Pharmacopeia XL / National Formulary 35</i> . Rockville, MD. US Pharmacopeial Convention, Inc. 2017: 5924.
7.	USP <795>. <i>United States Pharmacopeia XL / National Formulary 35</i> . Rockville, MD. US Pharmacopeial Convention, Inc. 2016: 675.

DISCLAIMER: MEDISCA NETWORK INC., HEREBY REFERRED TO AS 'THE NETWORK', HAS PROVIDED THE FORMULA AND INSTRUCTIONS ABOVE AS A MODEL FOR EDUCATIONAL PURPOSES ONLY ON THE BASIS OF THE RECOGNIZED COMPENDIA AND TEXTS REFERENCED AT THE END OF THIS DOCUMENT. THE NETWORK TAKES NO RESPONSIBILITY FOR THE VALIDITY, SCHEDULING OR ACCURACY OF THIS INFORMATION OR FOR ITS SAFETY OR EFFECTIVENESS, NOR FOR ANY USE THEREOF, WHICH IS AT THE SOLE RISK OF THE LICENSED PHARMACIST. ADJUSTMENTS MAY BE NEEDED TO MEET SPECIFIC PATIENT NEEDS AND IN ACCORDANCE WITH A LICENSED PRESCRIBER'S PRESCRIPTION. THE PHARMACIST MUST EMPLOY APPROPRIATE TESTS TO DETERMINE THE STABILITY OF THIS SUGGESTED FORMULA. THE NETWORK CANNOT BE HELD LIABLE TO ANY PERSON OR ENTITY CONCERNING CLAIMS, LOSS, OR DAMAGE CAUSED BY, OR ALLEGED TO BE CAUSED BY, DIRECTLY OR INDIRECTLY, THE USE OR MISUSE OF THE INFORMATION CONTAINED IN THIS SUGGESTED FORMULA. IN ALL CASES IT IS THE RESPONSIBILITY OF THE LICENSED PHARMACIST TO KNOW THE LAW, TO COMPOUND ANY FINISHED PRODUCT AND TO DISPENSE THESE PRODUCTS IN ACCORDANCE WITH FEDERAL AND STATE LAW. MEDISCA NETWORK INC. MAKES NO WARRANTIES WITH RESPECT TO INFRINGEMENT OR NON-INFRINGEMENT BY THE FORMULA OF ANY PATENT OR OTHER INTELLECTUAL PROPERTY OF ANY OTHER PARTY, AND IT IS THE RESPONSIBILITY OF THE PHARMACIST TO INVESTIGATE AND DETERMINE ANY SUCH ISSUE.