




MEDISCA® NETWORK INC.
TECHNICAL SUPPORT SERVICES
FORMULATION CHEMISTRY DEPARTMENT
TOLL-FREE: 866-333-7811
TELEPHONE: 514-905-5096
FAX: 514-905-5097
technicalservices@medisca.net

3/25/2008; page 1
TMP 045

Suggested Formula	Sodium Chloride 0.9% Intravenous or Intramuscular Injection (Solution, 100 mL)	FIN	F 002 239
-------------------	--	-----	-----------

SUGGESTED FORMULATION

Ingredient Listing	Qty.	Unit	NDC #	Supplier	Lot Number	Expiry Date
Sodium Chloride, USP	0.900	g				
Benzyl Alcohol, NF	1.0	mL				
Sterile Water For Injection, USP	90.0	mL		Ⓜ		
Sterile Water For Injection, USP	q.s to 100.0	mL				



DISCLAIMER: MEDISCA NETWORK INC. & RÉSEAU MEDISCA NETWORK INC., HEREBY REFERRED TO AS 'THE NETWORK', HAVE PROVIDED THE FORMULA AND INSTRUCTIONS ABOVE AS A MODEL FOR EDUCATIONAL PURPOSES ONLY ON THE BASIS OF THE RECOGNIZED COMPENDIA AND TEXTS REFERENCED AT THE END OF THIS DOCUMENT. THE NETWORK TAKES NO RESPONSIBILITY FOR THE VALIDITY OR ACCURACY OF THIS INFORMATION OR FOR ITS SAFETY OR EFFECTIVENESS, NOR FOR ANY USE THEREOF, WHICH IS AT THE SOLE RISK OF THE LICENSED PHARMACIST. THE PHARMACIST IS ADVISED THAT THE SUGGESTED FORMULA IS PRESENTED AT A SPECIFIED STRENGTH AND QUANTITY, AND THAT ADJUSTMENTS MAY BE NEEDED TO MEET SPECIFIC PATIENT NEEDS AND IN ACCORDANCE WITH A LICENSED PRESCRIBER'S PRESCRIPTION. FURTHER, THE FORMULA SHOULD ONLY BE PROVIDED UPON THE REQUEST AND AUTHORIZATION OF A LICENSED PRESCRIBER. PHARMACISTS USING THIS SUGGESTED FORMULA ARE ADVISED THAT ITS USE AS A MEANS OF THERAPEUTIC INTERVENTION IS BASED ON THE PRESCRIBER'S INDEPENDENT CLINICAL JUDGEMENT AND THE PROFESSIONAL EXPERTISE OF THE PHARMACIST. THE PHARMACIST MUST EMPLOY APPROPRIATE TESTS TO DETERMINE THE PRECISE STABILITY OF THIS SUGGESTED FORMULA. NO CLAIMS ARE BEING MADE AS TO THE SAFETY OR CLINICAL EFFICACY OF THIS SUGGESTED FORMULA. THE NETWORK CANNOT BE HELD LIABLE TO ANY PERSON OR ENTITY CONCERNING CLAIMS, LOSS, OR DAMAGE CAUSED BY, OR ALLEGED TO BE CAUSED BY, DIRECTLY OR INDIRECTLY, THE USE OR MISUSE OF THE INFORMATION CONTAINED IN THIS SUGGESTED FORMULA. FURTHERMORE, THE NETWORK, TAKES NO RESPONSIBILITY AND MAKES NO RECOMMENDATIONS IN DISPENSING THIS PRODUCT. IN ALL CASES IT IS THE RESPONSIBILITY OF THE LICENSED PHARMACIST TO KNOW THE LAW, TO COMPOUND ANY FINISHED PRODUCT AND TO DISPENSE THESE PRODUCTS IN ACCORDANCE WITH FEDERAL AND STATE LAW.



Suggested Formula	Sodium Chloride 0.9% Intravenous or Intramuscular Injection (Solution, 100 mL)	FIN	F 002 239
-------------------	--	-----	-----------

SPECIAL PREPARATORY CONSIDERATIONS

Ingredient-Specific Information

Light sensitive (protect from light whenever possible): Benzyl Alcohol

Hygroscopic (protect from moisture whenever possible): Sodium Chloride, Benzyl Alcohol

Suggested Preparatory Guidelines

Non-Sterile Preparation Sterile Preparation

Processing Error / Testing Considerations: To account for processing error, sterility and endotoxin testing considerations during preparation, it is suggested to measure an additional **5 to 9%** of the required quantities of ingredients.

Special Instruction: This formula must be prepared within the appropriate facilities under adequate environmental conditions, following the necessary guidelines and procedures as stated within *USP 797*. Only trained and qualified personnel must prepare this formula.

All heat stable, reusable materials and equipment must be sterilized and depyrogenated by dry heat sterilization at 250°C for 2 hours prior to use.

Every batch of final product compounded using this procedure must be sterility and endotoxin tested before being dispensed.

Protective apparel, such as a sterile gown, sterile gloves, shoe covers, head cap, eyewear and face-masks should always be worn. In addition, proper personnel cleansing must be done before entering the buffer or clean area.

Filter integrity must be validated by performing a filter stress test. If the test demonstrates that the filter might be defective, the solution must be discarded and remade.

This procedure requires the use of very small quantities of ingredients. All calculations and preparation techniques must be verified before dispensing the final product.

DISCLAIMER: MEDISCA NETWORK INC. & RÉSEAU MEDISCA NETWORK INC., HEREBY REFERRED TO AS 'THE NETWORK', HAVE PROVIDED THE FORMULA AND INSTRUCTIONS ABOVE AS A MODEL FOR EDUCATIONAL PURPOSES ONLY ON THE BASIS OF THE RECOGNIZED COMPENDIA AND TEXTS REFERENCED AT THE END OF THIS DOCUMENT. THE NETWORK TAKES NO RESPONSIBILITY FOR THE VALIDITY OR ACCURACY OF THIS INFORMATION OR FOR ITS SAFETY OR EFFECTIVENESS, NOR FOR ANY USE THEREOF, WHICH IS AT THE SOLE RISK OF THE LICENSED PHARMACIST. THE PHARMACIST IS ADVISED THAT THE SUGGESTED FORMULA IS PRESENTED AT A SPECIFIED STRENGTH AND QUANTITY, AND THAT ADJUSTMENTS MAY BE NEEDED TO MEET SPECIFIC PATIENT NEEDS AND IN ACCORDANCE WITH A LICENSED PRESCRIBER'S PRESCRIPTION. FURTHER, THE FORMULA SHOULD ONLY BE PROVIDED UPON THE REQUEST AND AUTHORIZATION OF A LICENSED PRESCRIBER. PHARMACISTS USING THIS SUGGESTED FORMULA ARE ADVISED THAT ITS USE AS A MEANS OF THERAPEUTIC INTERVENTION IS BASED ON THE PRESCRIBER'S INDEPENDENT CLINICAL JUDGEMENT AND THE PROFESSIONAL EXPERTISE OF THE PHARMACIST. THE PHARMACIST MUST EMPLOY APPROPRIATE TESTS TO DETERMINE THE PRECISE STABILITY OF THIS SUGGESTED FORMULA. NO CLAIMS ARE BEING MADE AS TO THE SAFETY OR CLINICAL EFFICACY OF THIS SUGGESTED FORMULA. THE NETWORK CANNOT BE HELD LIABLE TO ANY PERSON OR ENTITY CONCERNING CLAIMS, LOSS, OR DAMAGE CAUSED BY, OR ALLEGED TO BE CAUSED BY, DIRECTLY OR INDIRECTLY, THE USE OR MISUSE OF THE INFORMATION CONTAINED IN THIS SUGGESTED FORMULA. FURTHERMORE, THE NETWORK, TAKES NO RESPONSIBILITY AND MAKES NO RECOMMENDATIONS IN DISPENSING THIS PRODUCT. IN ALL CASES IT IS THE RESPONSIBILITY OF THE LICENSED PHARMACIST TO KNOW THE LAW, TO COMPOUND ANY FINISHED PRODUCT AND TO DISPENSE THESE PRODUCTS IN ACCORDANCE WITH FEDERAL AND STATE LAW.



Suggested Formula	Sodium Chloride 0.9% Intravenous or Intramuscular Injection (Solution, 100 mL)	FIN	F 002 239
-------------------	--	-----	-----------

SUGGESTED PREPARATION (for 100 mL)

Weigh and / or measure the following ingredients when appropriate:

Ingredient Listing	Qty.	Unit	Multiplication factor (*): ____	Processing Error	Qty. to measure
Sodium Chloride, USP §	0.900	g			
Benzyl Alcohol, NF §	1.0	mL			
Sterile Water For Injection, USP §	90.0	mL			
Sterile Water For Injection, USP §	q.s to 100.0	mL			

* Takes into account increased batch size conversions and density conversions, if required.

§ Weigh / measure just prior to use.

<u>Preparatory Instruction</u>	
IMPORTANT: All preparatory procedures must be performed using proper Aseptic Technique	
1.	<u>Equipment sterilization:</u> Following the manufacturer's specifications, sterilize and depyrogenate all heat stable, reusable materials and equipment, then return to ambient temperature.
2.	<u>Medium preparation:</u> A. Combine and mix the following ingredients together: - Benzyl Alcohol - Sterile Water For Injection (90.0 mL <i>plus</i> processing error adjustment) <u>Specifications:</u> Continuously mix. <u>End result:</u> Homogeneous liquid-like solution.

DISCLAIMER: MEDISCA NETWORK INC. & RÉSEAU MEDISCA NETWORK INC., HEREBY REFERRED TO AS 'THE NETWORK', HAVE PROVIDED THE FORMULA AND INSTRUCTIONS ABOVE AS A MODEL FOR EDUCATIONAL PURPOSES ONLY ON THE BASIS OF THE RECOGNIZED COMPENDIA AND TEXTS REFERENCED AT THE END OF THIS DOCUMENT. THE NETWORK TAKES NO RESPONSIBILITY FOR THE VALIDITY OR ACCURACY OF THIS INFORMATION OR FOR ITS SAFETY OR EFFECTIVENESS, NOR FOR ANY USE THEREOF, WHICH IS AT THE SOLE RISK OF THE LICENSED PHARMACIST. THE PHARMACIST IS ADVISED THAT THE SUGGESTED FORMULA IS PRESENTED AT A SPECIFIED STRENGTH AND QUANTITY, AND THAT ADJUSTMENTS MAY BE NEEDED TO MEET SPECIFIC PATIENT NEEDS AND IN ACCORDANCE WITH A LICENSED PRESCRIBER'S PRESCRIPTION. FURTHER, THE FORMULA SHOULD ONLY BE PROVIDED UPON THE REQUEST AND AUTHORIZATION OF A LICENSED PRESCRIBER. PHARMACISTS USING THIS SUGGESTED FORMULA ARE ADVISED THAT ITS USE AS A MEANS OF THERAPEUTIC INTERVENTION IS BASED ON THE PRESCRIBER'S INDEPENDENT CLINICAL JUDGEMENT AND THE PROFESSIONAL EXPERTISE OF THE PHARMACIST. THE PHARMACIST MUST EMPLOY APPROPRIATE TESTS TO DETERMINE THE PRECISE STABILITY OF THIS SUGGESTED FORMULA. NO CLAIMS ARE BEING MADE AS TO THE SAFETY OR CLINICAL EFFICACY OF THIS SUGGESTED FORMULA. THE NETWORK CANNOT BE HELD LIABLE TO ANY PERSON OR ENTITY CONCERNING CLAIMS, LOSS, OR DAMAGE CAUSED BY, OR ALLEGED TO BE CAUSED BY, DIRECTLY OR INDIRECTLY, THE USE OR MISUSE OF THE INFORMATION CONTAINED IN THIS SUGGESTED FORMULA. FURTHERMORE, THE NETWORK, TAKES NO RESPONSIBILITY AND MAKES NO RECOMMENDATIONS IN DISPENSING THIS PRODUCT. IN ALL CASES IT IS THE RESPONSIBILITY OF THE LICENSED PHARMACIST TO KNOW THE LAW, TO COMPOUND ANY FINISHED PRODUCT AND TO DISPENSE THESE PRODUCTS IN ACCORDANCE WITH FEDERAL AND STATE LAW.



Suggested Formula	Sodium Chloride 0.9% Intravenous or Intramuscular Injection (Solution, 100 mL)	FIN	F 002 239
3.	<p><u>Powder to medium integration:</u></p> <p>A. Incrementally add the Sodium Chloride to the following mixture: -Homogeneous liquid-like solution (Step 2A)</p> <p><u>Specifications:</u> Continuously mix, until all solid particles have completely dissolved.</p> <p><u>End result:</u> Homogeneous liquid-like solution.</p>		
4.	<p><u>Filling to volume:</u></p> <p>A. Add additional Sterile Water For Injection to the mixture (Step 3A) to fill to the required batch size (100.0 mL <i>plus</i> processing error adjustments).</p> <p><u>Specifications:</u> Continuously mix.</p> <p><u>End result:</u> Homogeneous liquid-like solution.</p>		
5.	<p><u>Filtering and transferring:</u></p> <p>Aseptically filter the solution through a 0.22-µm sterile filter into the recommended dispensing container (see Packaging requirements). Transfer the remainder into a separate dispensing container. This is to be used as the Test sample for sterility and endotoxin testing.</p>		
6.	<p><u>Filter integrity test:</u></p> <p>Validate filter integrity by performing a filter stress test. If the test demonstrates that the filter might be defective, the solution must be discarded and remade.</p>		
7.	<p><u>Sterility testing:</u></p> <p>Validate the Test sample for sterility and endotoxins, in accordance to current USP 797 regulatory guidelines.</p>		

DISCLAIMER: MEDISCA NETWORK INC. & RÉSEAU MEDISCA NETWORK INC., HEREBY REFERRED TO AS 'THE NETWORK', HAVE PROVIDED THE FORMULA AND INSTRUCTIONS ABOVE AS A MODEL FOR EDUCATIONAL PURPOSES ONLY ON THE BASIS OF THE RECOGNIZED COMPENDIA AND TEXTS REFERENCED AT THE END OF THIS DOCUMENT. THE NETWORK TAKES NO RESPONSIBILITY FOR THE VALIDITY OR ACCURACY OF THIS INFORMATION OR FOR ITS SAFETY OR EFFECTIVENESS, NOR FOR ANY USE THEREOF, WHICH IS AT THE SOLE RISK OF THE LICENSED PHARMACIST. THE PHARMACIST IS ADVISED THAT THE SUGGESTED FORMULA IS PRESENTED AT A SPECIFIED STRENGTH AND QUANTITY, AND THAT ADJUSTMENTS MAY BE NEEDED TO MEET SPECIFIC PATIENT NEEDS AND IN ACCORDANCE WITH A LICENSED PRESCRIBER'S PRESCRIPTION. FURTHER, THE FORMULA SHOULD ONLY BE PROVIDED UPON THE REQUEST AND AUTHORIZATION OF A LICENSED PRESCRIBER. PHARMACISTS USING THIS SUGGESTED FORMULA ARE ADVISED THAT ITS USE AS A MEANS OF THERAPEUTIC INTERVENTION IS BASED ON THE PRESCRIBER'S INDEPENDENT CLINICAL JUDGEMENT AND THE PROFESSIONAL EXPERTISE OF THE PHARMACIST. THE PHARMACIST MUST EMPLOY APPROPRIATE TESTS TO DETERMINE THE PRECISE STABILITY OF THIS SUGGESTED FORMULA. NO CLAIMS ARE BEING MADE AS TO THE SAFETY OR CLINICAL EFFICACY OF THIS SUGGESTED FORMULA. THE NETWORK CANNOT BE HELD LIABLE TO ANY PERSON OR ENTITY CONCERNING CLAIMS, LOSS, OR DAMAGE CAUSED BY, OR ALLEGED TO BE CAUSED BY, DIRECTLY OR INDIRECTLY, THE USE OR MISUSE OF THE INFORMATION CONTAINED IN THIS SUGGESTED FORMULA. FURTHERMORE, THE NETWORK, TAKES NO RESPONSIBILITY AND MAKES NO RECOMMENDATIONS IN DISPENSING THIS PRODUCT. IN ALL CASES IT IS THE RESPONSIBILITY OF THE LICENSED PHARMACIST TO KNOW THE LAW, TO COMPOUND ANY FINISHED PRODUCT AND TO DISPENSE THESE PRODUCTS IN ACCORDANCE WITH FEDERAL AND STATE LAW.



Suggested Formula	Sodium Chloride 0.9% Intravenous or Intramuscular Injection (Solution, 100 mL)	FIN	F 002 239
-------------------	--	-----	-----------

SUGGESTED PRESENTATION

Estimated Beyond-Use Date	14 days, refrigerated. BUD based on a successful sterility and endotoxin test result.	Packaging Requirements	Sterile, unit dose injection vials.
Auxiliary Labels	1 Use as directed. Do not exceed prescribed dose.	5	Keep refrigerated. Do not freeze.
	2 Discard container after use.	6	Consult your health care practitioner if any prescription or over-the-counter medications are currently being used or are prescribed for future use.
	3 Discard in the presence of particulate matter.	7	Do not use if discolored.
	4 Keep out of reach of children.		
Pharmacist Instructions	Add any auxiliary labels specific to the API to the dispensing container as deemed necessary.		
Patient Instructions	Contact your pharmacist in the event of adverse reactions.		

DISCLAIMER: MEDISCA NETWORK INC. & RÉSEAU MEDISCA NETWORK INC., HEREBY REFERRED TO AS 'THE NETWORK', HAVE PROVIDED THE FORMULA AND INSTRUCTIONS ABOVE AS A MODEL FOR EDUCATIONAL PURPOSES ONLY ON THE BASIS OF THE RECOGNIZED COMPENDIA AND TEXTS REFERENCED AT THE END OF THIS DOCUMENT. THE NETWORK TAKES NO RESPONSIBILITY FOR THE VALIDITY OR ACCURACY OF THIS INFORMATION OR FOR ITS SAFETY OR EFFECTIVENESS, NOR FOR ANY USE THEREOF, WHICH IS AT THE SOLE RISK OF THE LICENSED PHARMACIST. THE PHARMACIST IS ADVISED THAT THE SUGGESTED FORMULA IS PRESENTED AT A SPECIFIED STRENGTH AND QUANTITY, AND THAT ADJUSTMENTS MAY BE NEEDED TO MEET SPECIFIC PATIENT NEEDS AND IN ACCORDANCE WITH A LICENSED PRESCRIBER'S PRESCRIPTION. FURTHER, THE FORMULA SHOULD ONLY BE PROVIDED UPON THE REQUEST AND AUTHORIZATION OF A LICENSED PRESCRIBER. PHARMACISTS USING THIS SUGGESTED FORMULA ARE ADVISED THAT ITS USE AS A MEANS OF THERAPEUTIC INTERVENTION IS BASED ON THE PRESCRIBER'S INDEPENDENT CLINICAL JUDGEMENT AND THE PROFESSIONAL EXPERTISE OF THE PHARMACIST. THE PHARMACIST MUST EMPLOY APPROPRIATE TESTS TO DETERMINE THE PRECISE STABILITY OF THIS SUGGESTED FORMULA. NO CLAIMS ARE BEING MADE AS TO THE SAFETY OR CLINICAL EFFICACY OF THIS SUGGESTED FORMULA. THE NETWORK CANNOT BE HELD LIABLE TO ANY PERSON OR ENTITY CONCERNING CLAIMS, LOSS, OR DAMAGE CAUSED BY, OR ALLEGED TO BE CAUSED BY, DIRECTLY OR INDIRECTLY, THE USE OR MISUSE OF THE INFORMATION CONTAINED IN THIS SUGGESTED FORMULA. FURTHERMORE, THE NETWORK, TAKES NO RESPONSIBILITY AND MAKES NO RECOMMENDATIONS IN DISPENSING THIS PRODUCT. IN ALL CASES IT IS THE RESPONSIBILITY OF THE LICENSED PHARMACIST TO KNOW THE LAW, TO COMPOUND ANY FINISHED PRODUCT AND TO DISPENSE THESE PRODUCTS IN ACCORDANCE WITH FEDERAL AND STATE LAW.



Suggested Formula	Sodium Chloride 0.9% Intravenous or Intramuscular Injection (Solution, 100 mL)	FIN	F 002 239
-------------------	--	-----	-----------

REFERENCES

1.	USP <797> Pharmaceutical Compounding – Sterile Preparations. US Pharmacopeial Convention, Inc. <i>United States Pharmacopeia XXVIII / National Formulary 23</i> . Rockville, MD. 2004: 2461.
2.	Sodium Chloride. In: Rowe RC. <i>Handbook of Pharmaceutical Excipients, 4th Edition</i> . American Pharmaceutical Association; 2003: 556.
3.	Benzyl Alcohol. In: Rowe RC. <i>Handbook of Pharmaceutical Excipients, 4th Edition</i> . American Pharmaceutical Association; 2003: 53.
4.	Sodium Chloride (Monograph). <i>United States Pharmacopeia XXVIII / National Formulary 23</i> . Rockville, MD. US Pharmacopeial Convention, Inc. 2004: 2840.
5.	Benzyl Alcohol (Monograph). <i>United States Pharmacopeia XXVIII / National Formulary 23</i> . Rockville, MD. US Pharmacopeial Convention, Inc. 2004: 2965.

DISCLAIMER: MEDISCA NETWORK INC. & RÉSEAU MEDISCA NETWORK INC., HEREBY REFERRED TO AS 'THE NETWORK', HAVE PROVIDED THE FORMULA AND INSTRUCTIONS ABOVE AS A MODEL FOR EDUCATIONAL PURPOSES ONLY ON THE BASIS OF THE RECOGNIZED COMPENDIA AND TEXTS REFERENCED AT THE END OF THIS DOCUMENT. THE NETWORK TAKES NO RESPONSIBILITY FOR THE VALIDITY OR ACCURACY OF THIS INFORMATION OR FOR ITS SAFETY OR EFFECTIVENESS, NOR FOR ANY USE THEREOF, WHICH IS AT THE SOLE RISK OF THE LICENSED PHARMACIST. THE PHARMACIST IS ADVISED THAT THE SUGGESTED FORMULA IS PRESENTED AT A SPECIFIED STRENGTH AND QUANTITY, AND THAT ADJUSTMENTS MAY BE NEEDED TO MEET SPECIFIC PATIENT NEEDS AND IN ACCORDANCE WITH A LICENSED PRESCRIBER'S PRESCRIPTION. FURTHER, THE FORMULA SHOULD ONLY BE PROVIDED UPON THE REQUEST AND AUTHORIZATION OF A LICENSED PRESCRIBER. PHARMACISTS USING THIS SUGGESTED FORMULA ARE ADVISED THAT ITS USE AS A MEANS OF THERAPEUTIC INTERVENTION IS BASED ON THE PRESCRIBER'S INDEPENDENT CLINICAL JUDGEMENT AND THE PROFESSIONAL EXPERTISE OF THE PHARMACIST. THE PHARMACIST MUST EMPLOY APPROPRIATE TESTS TO DETERMINE THE PRECISE STABILITY OF THIS SUGGESTED FORMULA. NO CLAIMS ARE BEING MADE AS TO THE SAFETY OR CLINICAL EFFICACY OF THIS SUGGESTED FORMULA. THE NETWORK CANNOT BE HELD LIABLE TO ANY PERSON OR ENTITY CONCERNING CLAIMS, LOSS, OR DAMAGE CAUSED BY, OR ALLEGED TO BE CAUSED BY, DIRECTLY OR INDIRECTLY, THE USE OR MISUSE OF THE INFORMATION CONTAINED IN THIS SUGGESTED FORMULA. FURTHERMORE, THE NETWORK, TAKES NO RESPONSIBILITY AND MAKES NO RECOMMENDATIONS IN DISPENSING THIS PRODUCT. IN ALL CASES IT IS THE RESPONSIBILITY OF THE LICENSED PHARMACIST TO KNOW THE LAW, TO COMPOUND ANY FINISHED PRODUCT AND TO DISPENSE THESE PRODUCTS IN ACCORDANCE WITH FEDERAL AND STATE LAW.