



Suggested Formula	Fentanyl 50 µg/mL Epidural Injection (Solution, 5 mL)	FIN	F 004 217
-------------------	---	-----	-----------

SUGGESTED FORMULATION

Ingredient Listing	Qty.	Unit	NDC #	Supplier	Lot Number	Expiry Date
Fentanyl 50 µg/mL Injection (Sterile, Preservative Free) **	5.00	mL				

**Delivered as Fentanyl Citrate

SPECIAL PREPARATORY CONSIDERATIONS

Ingredient-Specific Information

Controlled substance (adhere to proper handling and documentation procedures)

Fentanyl Citrate

Light sensitive (protect from light whenever possible):

Fentanyl Citrate

Suggested Preparatory Guidelines

Non-Sterile Preparation Sterile Preparation

Processing Error / Testing Considerations: To account for processing error, sterility and endotoxin testing considerations during preparation, it is suggested to measure an additional **0 to 0%** of the required quantities of ingredients.

Special Instruction: This formula must be prepared within the appropriate facilities under adequate environmental conditions, following the necessary guidelines and procedures as stated within *USP 797*. Only trained and qualified personnel must prepare this formula.

All heat stable, reusable materials and equipment must be sterilized and depyrogenated by dry heat sterilization at 250°C for 2 hours prior to use.

Every batch of final product compounded using this procedure must be sterility and endotoxin tested before being dispensed.

Protective apparel, such as a sterile gown, sterile gloves, shoe covers, head cap, eyewear and face-masks should always be worn. In addition, proper personnel cleansing must be done before entering the buffer or clean area.

Filter integrity must be validated by performing a filter stress test. If the test demonstrates that the filter might be defective, the solution must be discarded and remade.

This procedure requires the use of very small quantities of ingredients. All calculations and preparation techniques must be verified before dispensing the final product.



Suggested Formula	Fentanyl 50 µg/mL Epidural Injection (Solution, 5 mL)	FIN	F 004 217
-------------------	---	-----	-----------

SUGGESTED PREPARATION (for 5 mL)

Weigh and / or measure the following ingredients when appropriate:

Ingredient Listing	Qty.	Unit	Multiplication factor (*): ____	Processing Error	Qty. to measure
Fentanyl 50 µg/mL Injection (Sterile, Preservative Free) §	5.00	mL			

* Takes into account increased batch size conversions and density conversions, if required.

§ Weigh / measure just prior to use.

<u>Preparatory Instruction</u>	
IMPORTANT: All preparatory procedures must be performed using proper Aseptic Technique	
1.	<u>Equipment sterilization:</u> Following the manufacturer's specifications, sterilize and depyrogenate all heat stable, reusable materials and equipment, then return to ambient temperature.
2.	<u>Medium integration:</u> Note: All manipulations must be done under a laminar airflow hood. Disinfect the commercial vials with Alcohol 70% prior to withdrawing the required amount of liquid. A. Measure and transfer the required amount of Fentanyl 50 µg/mL Injection (Sterile, Preservative Free).
3.	<u>Filtering and transferring:</u> Aseptically filter the solution through a 0.22-µm sterile filter into the recommended dispensing container (see Packaging requirements).
4.	<u>Filter integrity test:</u> Validate filter integrity by performing a filter stress test. If the test demonstrates that the filter might be defective, the solution must be discarded and remade.



Suggested Formula	Fentanyl 50 µg/mL Epidural Injection (Solution, 5 mL)	FIN	F 004 217
-------------------	---	-----	-----------

SUGGESTED PRESENTATION

Estimated Beyond-Use Date	14 days, refrigerated.	Packaging Requirements	Sterile, tight, light-resistant unit dose injection vials.	
Auxiliary Labels	1	Use as directed. Do not exceed prescribed dose.	8	For medical office use only.
	2	Keep out of reach of children.	9	Protect from light.
	3	Keep refrigerated. Do not freeze.	10	Controlled substance. Dangerous unless used as directed.
	4	Do not take with alcohol, sleep aids, tranquilizers or other CNS depressants.	11	May produce psychological and/or physical dependence.
	5	May impair mental and/or physical ability. Use care when operating a car or machinery.	12	Do not used if product changes color.
	6	Consult your health care practitioner if any other prescription or over-the-counter medications are currently being used or are prescribed for future use.	13	Preservative free solution, single use only. Discard any unused portion.
	7	Discard container after use.	14	Discard in the presence of particulate matter.
Pharmacist Instructions	Add any auxiliary labels specific to the active ingredients to the dispensing container as deemed necessary. IMPORTANT: TO BE ADMINISTERED ONLY BY THE PRESCRIBING PHYSICIAN.			
Patient Instructions	Contact your pharmacist in the event of adverse reactions.			



Suggested Formula	Fentanyl 50 µg/mL Epidural Injection (Solution, 5 mL)	FIN	F 004 217
-------------------	---	-----	-----------

REFERENCES

1.	USP <797>. <i>United States Pharmacopeia XXXII / National Formulary 27</i> . Rockville, MD. US Pharmacopeial Convention, Inc. 2009: 318.
2.	Parenteral Preparations. In: Allen, LV, Jr. <i>The Art, Science and Technology of Pharmaceutical Compounding Third Edition</i> . American Pharmaceutical Association; 2008: 313.
3.	Fentanyl Citrate. In: Sweetman SC, ed. <i>Martindale: The Complete Drug Reference, 34th Edition</i> . London, England: The Pharmaceutical Press; 2005: 40.
4.	Fentanyl (Monograph). In: O'Neil MJ. <i>The Merck Index 14th Edition</i> . Whitehouse Station, NJ: Merck & Co, Inc.; 2006: Monograph #4001.
5.	Fentanyl Citrate. In: Trissel LA. <i>Trissel's Stability of Compounded Formulations, 3rd Edition</i> . American Pharmaceutical Association; 2005: 172.
6.	Fentanyl Citrate (Monograph). <i>United States Pharmacopeia XXXII / National Formulary 27</i> . Rockville, MD. US Pharmacopeial Convention, Inc. 2009: 2358.
7.	Fentanyl Systemic. Thomson Micromedex. <i>USP DI – Drug Information for the Health Care Professional, 26th Edition</i> . Taunton, MA: US Pharmacopeial Convention, Inc; 2006: 1452.

DISCLAIMER: MEDISCA NETWORK INC. & RÉSEAU MEDISCA NETWORK INC., HEREBY REFERRED TO AS 'THE NETWORK', HAVE PROVIDED THE FORMULA AND INSTRUCTIONS ABOVE AS A MODEL FOR EDUCATIONAL PURPOSES ONLY ON THE BASIS OF THE RECOGNIZED COMPENDIA AND TEXTS REFERENCED AT THE END OF THIS DOCUMENT. THE NETWORK TAKES NO RESPONSIBILITY FOR THE VALIDITY OR ACCURACY OF THIS INFORMATION OR FOR ITS SAFETY OR EFFECTIVENESS, NOR FOR ANY USE THEREOF, WHICH IS AT THE SOLE RISK OF THE LICENSED PHARMACIST. ADJUSTMENTS MAY BE NEEDED TO MEET SPECIFIC PATIENT NEEDS AND IN ACCORDANCE WITH A LICENSED PRESCRIBER'S PRESCRIPTION. THE PHARMACIST MUST EMPLOY APPROPRIATE TESTS TO DETERMINE THE STABILITY OF THIS SUGGESTED FORMULA. THE NETWORK CANNOT BE HELD LIABLE TO ANY PERSON OR ENTITY CONCERNING CLAIMS, LOSS, OR DAMAGE CAUSED BY, OR ALLEGED TO BE CAUSED BY, DIRECTLY OR INDIRECTLY, THE USE OR MISUSE OF THE INFORMATION CONTAINED IN THIS SUGGESTED FORMULA. IN ALL CASES IT IS THE RESPONSIBILITY OF THE LICENSED PHARMACIST TO KNOW THE LAW, TO COMPOUND ANY FINISHED PRODUCT AND TO DISPENSE THESE PRODUCTS IN ACCORDANCE WITH FEDERAL AND STATE LAW.