



Suggested Formula	Lorazepam 2 mg/mL Intramuscular Injection (Solution, 100 mL)	FIN	F 004 989
-------------------	--	-----	-----------

### SUGGESTED FORMULATION

Ingredient Listing	Qty.	Unit	NDC #	Supplier	Lot Number	Expiry Date
Lorazepam, USP	0.200	g				
Polyethylene Glycol 400, NF	18.0	mL				
Benzyl Alcohol, NF	2.0	mL				
Propylene Glycol, USP	70.0	mL				
Propylene Glycol, USP	q.s. to 100.0	mL				

### SPECIAL PREPARATORY CONSIDERATIONS

#### Ingredient-Specific Information

**Light sensitive** (protect from light whenever possible):

*Lorazepam, Propylene Glycol, Benzyl Alcohol*

**Hygroscopic** (protect from moisture whenever possible):

*Propylene Glycol, Polyethylene Glycol 400*

**Controlled substance** (adhere to proper handling and documentation procedures)

*Lorazepam*

#### Suggested Preparatory Guidelines

Non-Sterile Preparation     Sterile Preparation

#### Processing Error / Testing Considerations:

To account for processing error, sterility and endotoxin testing considerations during preparation, it is suggested to measure an additional **5 to 9%** of the required quantities of ingredients.

#### Special Instruction:

This formula must be prepared within the appropriate facilities under adequate environmental conditions, following the necessary guidelines and procedures as stated within *USP 797*. Only trained and qualified personnel must prepare this formula.

All heat stable, reusable materials and equipment must be sterilized and depyrogenated by dry heat sterilization at 250°C for 2 hours prior to use.

Every batch of final product compounded using this procedure must be sterility and endotoxin tested before being dispensed.

Protective apparel, such as a sterile gown, sterile gloves, shoe covers, head cap, eyewear and face-masks should always be worn. In addition, proper personnel cleansing must be done before entering the buffer or clean area.

Filter integrity must be validated by performing a filter stress test. If the test demonstrates that the filter might be defective, the solution must be discarded and remade.

This procedure requires the use of very small quantities of ingredients. All calculations and preparation techniques must be verified before dispensing the final product.



Suggested Formula	Lorazepam 2 mg/mL Intramuscular Injection (Solution, 100 mL)	FIN	F 004 989
-------------------	--	-----	-----------

### SUGGESTED PREPARATION (for 100 mL)

Weigh and / or measure the following ingredients when appropriate:

Ingredient Listing	Qty.	Unit	Multiplication factor <sup>(*)</sup> : _____	Processing Error	Qty. to measure
Lorazepam, USP §	0.200	g			
Polyethylene Glycol 400, NF §	18.0	mL			
Benzyl Alcohol, NF §	2.0	mL			
Propylene Glycol, USP §	70.0	mL			
Propylene Glycol, USP §	q.s. to 100.0	mL			

\* Takes into account increased batch size conversions and density conversions, if required.

§ Weigh / measure just prior to use.

#### Preparatory Instruction

#### **IMPORTANT: All preparatory procedures must be performed using proper Aseptic Technique**

1. **Equipment sterilization:**

Following the manufacturer's specifications, sterilize and depyrogenate all heat stable, reusable materials and equipment, then return to ambient temperature.

2. **Medium preparation:**

A. Combine and mix the following ingredients together:

- Polyethylene Glycol 400
- Benzyl Alcohol
- Propylene Glycol (70.0 mL *plus* processing error adjustments).

End result: Homogeneous liquid-like solution.

3. **Medium Integration:**

A. Incrementally add the following ingredient to the Homogeneous liquid-like solution (Step 2A):

- Lorazepam

Specifications: Continuously mix until all solid particles have completely dissolved.

End result: Homogeneous liquid-like solution.



Suggested Formula	Lorazepam 2 mg/mL Intramuscular Injection (Solution, 100 mL)	FIN	F 004 989
-------------------	--	-----	-----------

4.	<p><b><u>Filling to volume:</u></b></p> <p>A. Add additional Propylene Glycol to the above mixture to fill to the required batch size (100.0 mL <i>plus</i> processing error adjustments).</p> <p><u>Specifications:</u> Continuously mix.</p> <p><u>End result:</u> Homogeneous liquid-like solution.</p>
5.	<p><b><u>Filtering and transferring:</u></b></p> <p>Using a water bath, warm the above solution to 60 – 70°C then aseptically filter the solution through a 0.22-µm sterile teflon filter into the recommended dispensing container (see Packaging requirements). Transfer the remainder into a separate dispensing container. This is to be used as the Test sample for sterility and endotoxin testing.</p>
6.	<p><b><u>Filter integrity test:</u></b></p> <p>Validate filter integrity by performing a filter stress test. If the test demonstrates that the filter might be defective, the solution must be discarded and remade.</p>
7.	<p><b><u>Sterility testing:</u></b></p> <p>Validate the Test sample for sterility and endotoxins, in accordance to current USP 797 regulatory guidelines.</p>



Suggested Formula	Lorazepam 2 mg/mL Intramuscular Injection (Solution, 100 mL)	FIN	F 004 989
-------------------	--	-----	-----------

**SUGGESTED PRESENTATION**

Estimated Beyond-Use Date	14 days, refrigerated, as per USP 797. BUD based on a successful sterility and endotoxin test result.	Packaging Requirements	Sterile, tightly closed, light-resistant injection vials.
Auxiliary Labels	1 Use as directed. Do not exceed prescribed dose.	8	Discard container after use.
	2 Keep out of reach of children.	9	Protect from light.
	3 Keep refrigerated. Do not freeze.	10	Equilibrate to room temperature before use.
	4 Consult your health care practitioner if any other prescription or over-the-counter medications are currently being used or are prescribed for future use.	11	Discard in the presence of particulate matter.
	5 Do not use if product changes color.	12	May impair mental and/or physical ability. Use care when operating a car or machinery.
	6 Do not take with alcohol, sleep aids, tranquilizers or other CNS depressants.	13	Keep in a dry place.
	7 May produce psychological and/or physical dependence.	14	Controlled substance. Dangerous unless used as directed.
Pharmacist Instructions	Add any auxiliary labels specific to the API to the dispensing container as deemed necessary.		
Patient Instructions	Contact your pharmacist in the event of adverse reactions.		



Suggested Formula	Lorazepam 2 mg/mL Intramuscular Injection (Solution, 100 mL)	FIN	F 004 989
-------------------	--	-----	-----------

## REFERENCES

1.	Parenteral Preparations. In: Allen, LV, Jr. <i>The Art, Science and Technology of Pharmaceutical Compounding Third Edition</i> . American Pharmaceutical Association; 2008: 313.
2.	Lorazepam Injection USP. In: Canadian Pharmacists Association. <i>Compendium of Pharmacists and Specialties, 2011</i> : 1390.
3.	Benzyl Alcohol. In: Rowe RC. <i>Handbook of Pharmaceutical Excipients, 6<sup>th</sup> Edition</i> . American Pharmaceutical Association; 2009: 64.
4.	Polyethylene Glycol. In: Rowe RC. <i>Handbook of Pharmaceutical Excipients, 6<sup>th</sup> Edition</i> . American Pharmaceutical Association; 2009: 517.
5.	Propylene Glycol. In: Rowe RC. <i>Handbook of Pharmaceutical Excipients, 6<sup>th</sup> Edition</i> . American Pharmaceutical Association; 2009: 592.
6.	Lorazepam. In: Sweetman SC, ed. <i>Martindale: The Complete Drug Reference, 36<sup>th</sup> Edition</i> . London, England: The Pharmaceutical Press; 2009: 1004.
7.	Lorazepam (Monograph). In: O'Neil MJ. <i>The Merck Index 14<sup>th</sup> Edition</i> . Whitehouse Station, NJ: Merck & Co, Inc.; 2006: Monograph #5579.
8.	Lorazepam. In: Trissel LA. <i>Trissel's Stability of Compounded Formulations, 4th Edition</i> . American Pharmaceutical Association; 2009: 342.
9.	Lorazepam (Monograph). <i>United States Pharmacopeia XXXIV / National Formulary 29</i> . Rockville, MD. US Pharmacopeial Convention, Inc. 2011: 3337.
10.	Lorazepam. Thomson Micromedex. <i>USP DI – Drug Information for the Health Care Professional, 26<sup>th</sup> Edition</i> . Taunton, MA: US Pharmacopeial Convention, Inc; 2006: 574.
11.	USP <797>. <i>United States Pharmacopeia XXXIV / National Formulary 29</i> . Rockville, MD. US Pharmacopeial Convention, Inc. 2011: 336.

DISCLAIMER: MEDISCA NETWORK INC. & RÉSEAU MEDISCA NETWORK INC., HEREBY REFERRED TO AS 'THE NETWORK', HAVE PROVIDED THE FORMULA AND INSTRUCTIONS ABOVE AS A MODEL FOR EDUCATIONAL PURPOSES ONLY ON THE BASIS OF THE RECOGNIZED COMPENDIA AND TEXTS REFERENCED AT THE END OF THIS DOCUMENT. THE NETWORK TAKES NO RESPONSIBILITY FOR THE VALIDITY OR ACCURACY OF THIS INFORMATION OR FOR ITS SAFETY OR EFFECTIVENESS, NOR FOR ANY USE THEREOF, WHICH IS AT THE SOLE RISK OF THE LICENSED PHARMACIST. ADJUSTMENTS MAY BE NEEDED TO MEET SPECIFIC PATIENT NEEDS AND IN ACCORDANCE WITH A LICENSED PRESCRIBER'S PRESCRIPTION. THE PHARMACIST MUST EMPLOY APPROPRIATE TESTS TO DETERMINE THE STABILITY OF THIS SUGGESTED FORMULA. THE NETWORK CANNOT BE HELD LIABLE TO ANY PERSON OR ENTITY CONCERNING CLAIMS, LOSS, OR DAMAGE CAUSED BY, OR ALLEGED TO BE CAUSED BY, DIRECTLY OR INDIRECTLY, THE USE OR MISUSE OF THE INFORMATION CONTAINED IN THIS SUGGESTED FORMULA. IN ALL CASES IT IS THE RESPONSIBILITY OF THE LICENSED PHARMACIST TO KNOW THE LAW, TO COMPOUND ANY FINISHED PRODUCT AND TO DISPENSE THESE PRODUCTS IN ACCORDANCE WITH FEDERAL AND STATE LAW.