

3/19/2012; page 1

Suggested Formula	Midazolam 5 mg/mL Intravenous Injection (Solution, 10 mL)	FIN	F 004 993v2
----------------------	---	-----	-------------

# SUGGESTED FORMULATION

Ingredient Listing	Qty.	Unit	NDC #	Supplier	Lot Number	Expiry Date
Midazolam, USP	0.050	g				
Sodium Chloride, USP	0.08	g				
Edetate Disodium 0.2% Stock Solution †	0.5	mL				
Benzyl Alcohol, NF	0.1	mL				
Sterile Water for Injection, USP	8.0	mL				
Sterile Water for Injection, USP	q.s. to 10.0	mL	$\odot$			
Hydrochloric Acid 1N solution	As required			1		
			JY X	)		
† Edetate Disodium 0.2% Stock Solution		S		4		
Edetate Disodium, USP	0.02	g				
Sterile Water for Injection, USP	10.0	mL				



3/19/2012; page 2

	Suggested FormulaMidazolam 5 mg/mL Intravenous Injection (Solution, 10 mL)					
SP		PARATORY CONSIL	DERATIONS			
	Ingredient-	Specific Information				
	Light se	<b>nsitive</b> (protect from lig	th whenever possible): Midazolam			
		<b>led substance</b> (adhere t ntation procedures)	o proper handling and Midazolam			
	Hygroso	c <b>opic</b> (protect from mois	sture whenever possible): Edetate Disodium			
	Suggested 1	Preparatory Guidelines	8			
	Γ	Non-Sterile Preparati	on Sterile Preparation			
		ocessing Error / esting Considerations:	To account for processing error, pH testing, sterility and considerations during preparation, it is suggested to measure an ac of the required quantities of ingredients.			
	<u>Sp</u>	ecial Instruction:	This formula must be prepared within the appropriate facilities under environmental conditions, following the necessary guidelines and p within <i>USP 797</i> . Only trained and qualified personnel must prepare	rocedure	es as stated	
			All heat stable, reusable materials and equipment must be sterilized by dry heat sterilization at 250°C for 2 hours prior to use.	and dep	byrogenated	
			Every batch of final product compounded using this procedure mus endotoxin tested before being dispensed.	be ster	ility and	
			Protective apparel, such as a sterile gown, sterile gloves, shoe cover eyewear and face-masks should always be worn. In addition, proper cleansing must be done before entering the buffer or clean area.			
			Filter integrity must be validated by performing a filter stress test. I demonstrates that the filter might be defective, the solution must be remade.			
			This procedure requires the use of very small quantities of ingredien and preparation techniques must be verified before dispensing the f			



3/19/2012; page 3

Suggested Formula	Midazolam 5 mg/mL Intravenous Injection (Solution, 10 mL)	FIN	F 004 993v2	
----------------------	---	-----	-------------	--

# SUGGESTED PREPARATION (for 10 mL)

Weigh and / or measure the following ingredients when appropriate:

Ingredient Listing	Qty.	Unit	Multiplication factor <sup>(*)</sup> :	Processing Error	Qty. to measure
Midazolam, USP §	0.050	g			
Sodium Chloride, USP §	0.08	g			
Edetate Disodium 0.2% Stock Solution † §	0.5	mL			
Benzyl Alcohol, NF §	0.1	mL	(C)		
Sterile Water for Injection, USP §	8.0	mL			
Sterile Water for Injection, USP §	q.s. to 10.0	mL			
Hydrochloric Acid 1N solution §	As required		1		
† Edetate Disodium 0.2% Stock Solution					
Edetate Disodium, USP §	0.02	g			
Sterile Water for Injection, USP §	10.0	mL			

\* Takes into account increased batch size conversions and density conversions, if required.

§ Weigh / measure just prior to use.

## **Preparatory Instruction**

IMPORTANT: All preparatory procedures must be performed using proper Aseptic Technique

1. **Equipment sterilization:** 

Following the manufacturer's specifications, sterilize and depyrogenate all heat stable, reusable materials and equipment, then return to ambient temperature.

2. † Edetate Disodium 0.2% Stock Solution preparation:

A. Incrementally add the Edetate Disodium to the Sterile Water for Injection (10.0 mL) and mix until dissolved.



3/19/2012; page 4

Suggested Formula		Midazolam 5 mg/mL Intravenous Injection (Solution, 10 mL)	FIN	F 004 993v2				
3.	Powder-liquid preparation:							
		the given order, sequentially add the following ingredients to the Sterile Water for Injection rocessing error adjustments).	(8.0 m	L plus				
	<ul> <li>-Benzyl Alcohol</li> <li>-Sodium Chloride</li> <li>-Edetate Disodium 0.2% Stock Solution (0.5 mL <i>plus</i> processing error adjustment)</li> <li>-Midazolam</li> </ul>							
	<u>S</u>	pecifications: Continuously mix until all solid particles have completely dispersed.						
	E	nd result: Homogeneous liquid-like dispersion.						
	N	tote: Add the next ingredient, once the previous one has been completely added and dispersed	1.					
4.	<u>pH te</u>	sting:						
	A. C	braw an appropriate amount of the mixture (Step 2A).						
	В. Т	est the pH of the sample. It should lie between 3.1 and 3.5						
	C. <u>I</u>	the pH > 3.5, carefully add, in a dropwise fashion, the Hydrochloric Acid 1N solution to the	mixtur	<u>e:</u>				
	<ol> <li>Draw and transfer 1 or 2 drops of the Hydrochloric Acid 1N solution to the mixture.</li> <li>Stir for at least 5 minutes to evenly disperse the Hydrochloric Acid 1N solution.</li> <li>Re-test the pH.</li> <li>Continue to add the Hydrochloric Acid 1N solution until all solid particles have completely dissolved and a pH of 3.1 to 3.5 is obtained.</li> </ol>							
	IMPORTANT: Do not allow the pH to fall below 3.1							
5.	Fillin	g to volume:						
		dd additional Sterile Water for Injection to the above mixture to fill to the required batch size rocessing error adjustments).	(10.0	mL <i>plus</i>				
	<u>S</u>	pecification: Continuously mix						
	<u>E</u>	nd result: Homogeneous liquid-like solution						
6.	Filter	ing and transferring:						
		ically filter the required amount of solution through a $0.22$ -µm sterile filter into the recominers (see Packaging requirements) and sample containers for sterility and endotoxin testing.	nmende	ed dispensing				
7.	Filter	· integrity test:						
		ate filter integrity by performing a filter stress test. If the test demonstrates that the filter migh on must be discarded and remade.	t be de	fective, the				
8.	Steri	ity testing:						
	Valid	ate the Test samples for sterility and endotoxins, in accordance to current USP 797 regulatory	guide	ines.				



3/19/2012; page 5

Suggested Formula Midazolam 5 mg/mL Intravenous Injection (Solution, 10 mL)

FIN F 004 993v2

SUGGESTED PR	ESE	NTATION			
Estimated Beyond-Use Date		<ul><li>14 days, refrigerated as per USP 797.</li><li>BUD based on a successful sterility and endotoxin test result.</li></ul>	Packaging Requirements		Sterile, tightly closed, light-resistant unit dose injection vials.
	1	Use as directed. Do not exceed dose.	l prescribed	8	Do not used if product changes color.
	2	Keep out of reach of children.		9	Protect from light.
	3	Keep refrigerated. Do not freeze		10	For medical office use only.
	4	Do not take with alcohol, tranquilizers or other CNS depre		11	May produce psychological and/or physical dependence.
Auxiliary Labels	5	May impair mental and/or phys Use care when operating machinery.		12	Controlled substance. Dangerous unless used as directed.
	6	Consult your health care practit other prescription or over- medications are currently being prescribed for future use.	-the-counter	13	Administered slowly via intravenous injection in incremental doses or intravenous infusion. It may be diluted with sodium chloride 0.9% or dextrose 5% to facilitate slow administration.
	7	Discard in the presence of matter.	particulate	14	Equilibrate to room temperature before use.
Pharmacist Instructions	Add any auxiliary labels specific to the active ingredients to the dispensing container as deemed necessary.				
Patient Instructions	Contact your pharmacist in the event of adverse reactions.				

## REFERENCES

1.	Parenteral Preparations. In: Allen, LV, Jr. <i>The Art, Science and Technology of Pharmaceutical Compounding Third Edition</i> . American Pharmaceutical Association; 2008: 313.
2.	Midazolam. In: Sweetman SC, ed. <i>Martindale: The Complete Drug Reference, 36<sup>th</sup> Edition</i> . London, England: The Pharmaceutical Press; 2009: 1007.
3.	Midazolam (Monograph). In: O'Neil MJ. <i>The Merck Index 14<sup>th</sup> Edition</i> . Whitehouse Station, NJ: Merck & Co, Inc.; 2006: Monograph #6178.
4.	Midazolam (Monograph). <i>United States Pharmacopeia XXXIV / National Formulary 29</i> . Rockville, MD. US Pharmacopeial Convention, Inc. 2011: 3530.
5.	USP <797>. United States Pharmacopeia XXXIV / National Formulary 29. Rockville, MD. US Pharmacopeial Convention, Inc. 2011: 336.

DISCLAIMER: MEDISCA NETWORK INC. & RÉSEAU MEDISCA NETWORK INC., HEREBY REFERRED TO AS 'THE NETWORK', HAVE PROVIDED THE FORMULA AND INSTRUCTIONS ABOVE AS A MODEL FOR EDUCATIONAL PURPOSES ONLY ON THE BASIS OF THE RECOGNIZED COMPENDIA AND TEXTS REFERENCED AT THE END OF THIS DOCUMENT. THE NETWORK TAKES NO RESPONSIBILITY FOR THE VALIDITY OR ACCURACY OF THIS INFORMATION OR FOR ITS SAFETY OR EFFECTIVENESS, NOR FOR ANY USE THEREOF, WHICH IS AT THE SOLE RISK OF THE LICENSED PHARMACIST. ADJUSTMENTS MAY BE NEEDED TO MEET SPECIFIC PATIENT NEEDS AND IN ACCORDANCE WITH A LICENSED PRESCRIBER'S PRESCRIPTION. THE PHARMACIST MUST EMPLOY APPROPRIATE TESTS TO DETERMINE THE STABILITY OF THIS SUGGESTED FORMULA. THE NETWORK CANNOT BE HELD LIABLE TO ANY PERSON OR ENTITY CONCERNING CLAIMS, LOSS, OR DAMAGE CAUSED BY, OR ALLEGED TO BE CAUSED BY, DIRECTLY OR INDIRECTLY, THE USE OR MISUSE OF THE INFORMATION CONTAINED IN THIS SUGGESTED FORMULA. IN ALL CASES IT IS THE RESPONSIBILITY OF THE LICENSED PHARMACIST TO KNOW THE LAW, TO COMPOUND ANY FINISHED PRODUCT AND TO DISPENSE THESE PRODUCTS IN ACCORDANCE WITH FEDERAL AND STATE LAW.