



Suggested Formula	Fluvoxamine Maleate 50 mg/5 mL Oral Liquid (Suspension, 100 mL)	FIN	F 002 593v2
-------------------	---	-----	-------------

SUGGESTED FORMULATION

Ingredient Listing	Qty.	Unit	NDC #	Supplier	Lot Number	Expiry Date
Fluvoxamine Maleate, BP	1.000	g				
Glycerin, USP	5.0	mL				
Medisca Oral Suspend (Suspending Vehicle)	50.0	mL				
Medisca Oral Syrup (Flavored Vehicle)	q.s. to 100.0	mL				

SPECIAL PREPARATORY CONSIDERATIONS

Ingredient-Specific Information

Hygroscopic (protect from moisture whenever possible):

Glycerin

Light Sensitive (protect from light whenever possible):

Fluvoxamine Maleate

Suggested Preparatory Guidelines

Non-Sterile Preparation Sterile Preparation

Processing Error /

Testing Considerations:

To account for processing error considerations during preparation, it is suggested to measure an additional **5 to 9%** of the required quantities of ingredients.

Special Instruction:

Protective apparel, such as a lab coat, disposable gloves, eyewear and face-masks should always be worn.

This procedure requires the use of very small quantities of ingredients. All calculations and preparation techniques must be verified before dispensing the final product.



Suggested Formula	Fluvoxamine Maleate 50 mg/5 mL Oral Liquid (Suspension, 100 mL)	FIN	F 002 593v2
-------------------	---	-----	-------------

SUGGESTED PREPARATION (for 100 mL)

Weigh and / or measure the following ingredients when appropriate:

Ingredient Listing	Qty.	Unit	Multiplication factor (*): ____	Processing Error	Qty. to measure
Fluvoxamine Maleate, BP §	1.000	g			
Glycerin, USP §	5.0	mL			
Medisca Oral Suspend (Suspending Vehicle)	50.0	mL			
Medisca Oral Syrup (Flavored Vehicle)	q.s. to 100.0	mL			

§ Weigh / measure just prior to use.

* Takes into account increased batch size conversions and density conversions, if required.

Preparatory Instruction	
1.	<p><u>Powder-liquid preparation:</u></p> <p>A. Triturate the Fluvoxamine Maleate to form a fine, homogeneous powder.</p> <p>B. Levigate the fine, homogeneous powder (Step 1A) with the Glycerin.</p> <p><u>End result:</u> Homogeneous liquid-like dispersion.</p>
2.	<p><u>Medium integration:</u></p> <p>A. Incrementally add the homogeneous liquid-like dispersion (Step 1B) to the Oral Suspend (Suspending Vehicle).</p> <p><u>Specifications:</u> Continuously mix.</p> <p><u>End result:</u> Homogeneous liquid-like dispersion.</p>
3.	<p><u>Filling to volume:</u></p> <p>A. Add Oral Syrup (Flavored Vehicle) to the mixture (Step 2A) to fill to the required batch size (100.0 mL <i>plus</i> processing error adjustments).</p> <p><u>Specifications:</u> Continuously mix.</p> <p><u>End result:</u> Homogeneous liquid-like dispersion.</p>



Suggested Formula	Fluvoxamine Maleate 50 mg/5 mL Oral Liquid (Suspension, 100 mL)	FIN	F 002 593v2
-------------------	---	-----	-------------

4.	<p><u>Product transfer:</u></p> <p>A. Transfer the final product into the specified dispensing container (see “Packaging requirements”).</p> <p><u>Note:</u> Continuously mix the final product during the transfer process into the recommended dispensing containers in order to maintain homogeneity.</p>
----	---

SUGGESTED PRESENTATION

Estimated Beyond-Use Date		Packaging Requirements	
	14 days, refrigerated, as per USP.		- Tightly closed, light-resistant dispensing bottle. - To be administered with a metered dose-measuring device.
Auxiliary Labels	1	Use as directed. Do not exceed prescribed dose.	6 Cap tightly after use.
	2	Keep out of reach of children.	7 Shake well before use.
	3	Consult your health care practitioner if any other prescription or over-the-counter medications are currently being used or are prescribed for future use.	8 Keep refrigerated. Do not freeze.
	4	Do not take with alcohol, sleep aids, tranquilizers or other CNS depressants.	9 Protect from light.
	5	May impair mental and/or physical ability. Use care when operating a car or machinery.	
Pharmacist Instructions	Add any auxiliary labels specific to the active to the dispensing container as deemed necessary.		
Patient Instructions	Contact your pharmacist in the event of adverse reactions.		



Suggested Formula	Fluvoxamine Maleate 50 mg/5 mL Oral Liquid (Suspension, 100 mL)	FIN	F 002 593v2
-------------------	---	-----	-------------

REFERENCES

1.	USP <795>. <i>United States Pharmacopeia XXVIII / National Formulary 23</i> . Rockville, MD. US Pharmacopeial Convention, Inc. 2004: 2457.
2.	Glycerin. In: Rowe RC. <i>Handbook of Pharmaceutical Excipients, 4th Edition</i> . American Pharmaceutical Association; 2003: 257.
3.	Fluvoxamine Maleate. In: Sweetman SC, ed. <i>Martindale: The Complete Drug Reference, 34th Edition</i> . London, England: The Pharmaceutical Press; 2005: 298.
4.	Fluvoxamine (Monograph). In: O'Neil MJ. <i>The Merck Index 14th Edition</i> . Whitehouse Station, NJ: Merck & Co, Inc.; 2006: #4219.
5.	Fluvoxamine. Thomson Micromedex. <i>USP DI – Drug Information for the Health Care Professional, 26th Edition</i> . Taunton, MA: US Pharmacopeial Convention, Inc; 2006: 1533.

DISCLAIMER: MEDISCA NETWORK INC., HEREBY REFERRED TO AS 'THE NETWORK', HAS PROVIDED THE FORMULA AND INSTRUCTIONS ABOVE AS A MODEL FOR EDUCATIONAL PURPOSES ONLY ON THE BASIS OF THE RECOGNIZED COMPENDIA AND TEXTS REFERENCED AT THE END OF THIS DOCUMENT. THE NETWORK TAKES NO RESPONSIBILITY FOR THE VALIDITY OR ACCURACY OF THIS INFORMATION OR FOR ITS SAFETY OR EFFECTIVENESS, NOR FOR ANY USE THEREOF, WHICH IS AT THE SOLE RISK OF THE LICENSED PHARMACIST. ADJUSTMENTS MAY BE NEEDED TO MEET SPECIFIC PATIENT NEEDS AND IN ACCORDANCE WITH A LICENSED PRESCRIBER'S PRESCRIPTION. THE PHARMACIST MUST EMPLOY APPROPRIATE TESTS TO DETERMINE THE STABILITY OF THIS SUGGESTED FORMULA. THE NETWORK CANNOT BE HELD LIABLE TO ANY PERSON OR ENTITY CONCERNING CLAIMS, LOSS, OR DAMAGE CAUSED BY, OR ALLEGED TO BE CAUSED BY, DIRECTLY OR INDIRECTLY, THE USE OR MISUSE OF THE INFORMATION CONTAINED IN THIS SUGGESTED FORMULA. IN ALL CASES IT IS THE RESPONSIBILITY OF THE LICENSED PHARMACIST TO KNOW THE LAW, TO COMPOUND ANY FINISHED PRODUCT AND TO DISPENSE THESE PRODUCTS IN ACCORDANCE WITH FEDERAL AND STATE LAW.