



| | | | |
|-------------------|---|-----|-----------|
| Suggested Formula | Nystatin 100,000 IU/mL Oral Liquid (Suspension, 100 mL) | FIN | F 008 096 |
|-------------------|---|-----|-----------|

SUGGESTED FORMULATION

| Ingredient Listing | Qty. | Unit | NDC # | Supplier | Lot Number | Expiry Date |
|---|---------------|------|-------|----------|------------|-------------|
| Nystatin, USP | 10 | MU | | | | |
| Glycerin, USP | 5.0 | mL | | | | |
| Xanthan Gum, NF | 1.5 | g | | | | |
| Medisca Oral Syrup (Flavored Syrup Vehicle) | 50.0 | mL | | | | |
| Medisca Oral Syrup (Flavored Syrup Vehicle) | q.s. to 100.0 | mL | | | | |
| Sodium Hydroxide 10% Solution | As required | | | | | |
| Citric Acid 10% Solution | As required | | | | | |

SPECIAL PREPARATORY CONSIDERATIONS

Ingredient-Specific Information

| | |
|--|-----------------------|
| Light Sensitive (protect from light whenever possible): | Nystatin |
| Hygroscopic (protect from moisture whenever possible): | Glycerin, Nystatin |
| Oxygen Sensitive (protect from oxygen whenever possible): | Nystatin |
| Moisture Sensitive (protect from humidity whenever possible): | Nystatin, Citric Acid |
| Air Sensitive (protect from air whenever possible): | Nystatin |
| Heat Sensitive (protect from heat whenever possible): | Nystatin |



| | | | |
|-------------------|---|-----|-----------|
| Suggested Formula | Nystatin 100,000 IU/mL Oral Liquid (Suspension, 100 mL) | FIN | F 008 096 |
|-------------------|---|-----|-----------|

SPECIAL PREPARATORY CONSIDERATIONS (CONTINUED)

Suggested Preparatory Guidelines

Non-Sterile Preparation Sterile Preparation

Processing Error / Testing Considerations: To account for processing error and pH testing considerations during preparation, it is suggested to measure an additional **1 to 3%** of the required quantities of ingredients.

Special Instruction: This formula may contain one or more Active Pharmaceutical Ingredients (APIs) that may be classified as hazardous, please refer & verify the current NIOSH list of Antineoplastic and Other Hazardous Drugs in Healthcare Settings, 2016. **General Chapter <800> Hazardous Drugs – Handling in Healthcare Settings** was formally published February 1, 2016 in the First Supplement to USP 39-NF 34 and has a delayed **official implementation date of December 31st, 2019**.

This formula must be prepared within the appropriate facilities under adequate environmental conditions, following the necessary guidelines and procedures as stated within *USP 795* and *USP 800*, when handling hazardous drugs. Only trained and qualified personnel must prepare this formula.

All required personal protective equipment (hazardous if applicable), such as but not limited to, lab coat, protective sleeves, gloves both inner and outer if applicable, dedicated shoe covers, hairnet, beard cover, eyewear, appropriate face mask, respirator and face shield, etc., where applicable must be worn at all times.

If applicable, follow all required procedures for hazardous drug handling including but not limited to procurement, transport, storage, preparation, dispensing, administration, clean up (spills) & disposal.

If you are a registered 503B facility, please refer to all relevant guidance documents including but not limited to the Code of Federal Regulations (CFR), Guidance for Industry (GFIs) and Compliance Policy Guides (CPGs).

This procedure requires the use of very small quantities of ingredients. All calculations and preparation techniques must be verified before dispensing the final product.



| | | | |
|-------------------|---|-----|-----------|
| Suggested Formula | Nystatin 100,000 IU/mL Oral Liquid (Suspension, 100 mL) | FIN | F 008 096 |
|-------------------|---|-----|-----------|

SUGGESTED PREPARATION (for 100 mL)

Weigh and / or measure the following ingredients when appropriate:

| Ingredient Listing | Qty. | Unit | Multiplication factor (*): _____ | Processing Error | Qty. to measure |
|---|---------------|------|----------------------------------|------------------|-----------------|
| Nystatin, USP § | 10 | MU | | | |
| Glycerin, USP § | 5.0 | mL | | | |
| Xanthan Gum, NF | 1.5 | g | | | |
| Medisca Oral Syrup (Flavored Syrup Vehicle) | 50.0 | mL | | | |
| Medisca Oral Syrup (Flavored Syrup Vehicle) | q.s. to 100.0 | mL | | | |
| Sodium Hydroxide 10% Solution | As required | | | | |
| Citric Acid 10% Solution | As required | | | | |

* Takes into account increased batch size conversions and density conversions, if required.

§ Weigh / measure just prior to use.





| | | | |
|-------------------|---|-----|-----------|
| Suggested Formula | Nystatin 100,000 IU/mL Oral Liquid (Suspension, 100 mL) | FIN | F 008 096 |
|-------------------|---|-----|-----------|

Preparatory Instruction

1. Ingredient quantification:

A. Determine the quantity (in g) of Nystatin required to make a Nystatin 10 MU Oral Liquid, batch size (100 mL)

| | |
|--|---------------|
| Quantity of Nystatin (in Units) required | 10,000,000 IU |
| DIVIDED BY | |
| Nystatin biopotency assay result (from Certificate of Analysis) | _____ IU/mg |
| EQUALS | |
| i. Quantity of Nystatin (in milligrams) required | _____ mg |
| MULTIPLIED BY | |
| Multiplication factor – milligrams to grams | 0.001 |
| EQUALS | |
| ii. Quantity of Nystatin (in grams) required | _____ g |
| MULTIPLIED BY | |
| Processing error adjustments (5 to 9%) | 1.05 to 1.09 |
| EQUALS | |
| iii. Quantity of Nystatin needed <i>plus</i> processing error adjustments | _____ g |

2. Powder-liquid preparation:

A. Levigate the Nystatin (amount determined in Step 1Aiii) with the Glycerin.

End result: Homogeneous liquid-like dispersion.



| | | | |
|-------------------|---|-----|-----------|
| Suggested Formula | Nystatin 100,000 IU/mL Oral Liquid (Suspension, 100 mL) | FIN | F 008 096 |
|-------------------|---|-----|-----------|

| | |
|----|---|
| 3. | <p><u>Medium integration:</u></p> <p>A. In the given order, sequentially add the following ingredients to the Medisca Oral Syrup (Flavored Syrup Vehicle) (50.0 mL <i>plus</i> processing error adjustments):</p> <ul style="list-style-type: none">-Homogeneous liquid-like dispersion (Step 2A)-Xanthan Gum <p><u>Specifications:</u> Continuously mix, using high-shear mixing techniques.</p> <p><u>End result:</u> Homogeneous liquid-like dispersion.</p> <p><u>Note:</u> Add the next ingredient, once the previous one has been completely added and dispersed.</p> |
| 4. | <p><u>Filling to volume:</u></p> <p>A. Add additional Medisca Oral Syrup (Flavored Syrup Vehicle) to the mixture (Step 3A) to fill to the required batch size (100.0 mL <i>plus</i> processing error adjustments).</p> <p><u>Specifications:</u> Continuously mix, using high-shear mixing techniques.</p> <p><u>End result:</u> Homogeneous liquid-like dispersion.</p> |
| 5. | <p><u>pH testing:</u></p> <p>A. Draw an appropriate amount of the mixture (Step 4A).</p> <p>B. Test the pH of the sample. It should lie between 6.0 and 7.0.</p> <p>C. <u>If the pH < 6.0, carefully add, in a dropwise fashion, the Sodium Hydroxide 10% Solution to the mixture:</u></p> <ol style="list-style-type: none">1. Draw and transfer 1 or 2 drops of the Sodium Hydroxide 10% Solution to the mixture.2. Stir for at least 5 minutes to evenly disperse the Sodium Hydroxide 10% Solution.3. Re-test the pH.4. Continue to add the Sodium Hydroxide 10% Solution until the pH of 6.0 to 7.0 is obtained. <p>IMPORTANT: Do not allow the pH to rise above 7.0.</p> <p>D. <u>If the pH > 7.0, carefully add, in a dropwise fashion, the Citric Acid 10% Solution to the mixture:</u></p> <ol style="list-style-type: none">1. Draw and transfer 1 or 2 drops of the Citric Acid 10% Solution to the mixture.2. Stir for at least 5 minutes to evenly disperse the Citric Acid 10% Solution.3. Re-test the pH.4. Continue to add the Citric Acid 10% Solution until the pH of 6.0 to 7.0 is obtained. <p>IMPORTANT: Do not allow the pH to fall below 6.0.</p> |



| | | | |
|-------------------|---|-----|-----------|
| Suggested Formula | Nystatin 100,000 IU/mL Oral Liquid (Suspension, 100 mL) | FIN | F 008 096 |
|-------------------|---|-----|-----------|

| | |
|----|---|
| 6. | <p><u>Product transfer:</u></p> <p>A. Transfer the final product into the specified dispensing container (see “Packaging requirements”).</p> <p><u>Note:</u> Continuously mix the final product during the transfer process into the recommended dispensing containers in order to maintain homogeneity.</p> |
|----|---|

SUGGESTED PRESENTATION

| | | | |
|---------------------------|---|------------------------|--|
| Estimated Beyond-Use Date | 14 days, refrigerated, as per USP. | Packaging Requirements | - Tightly closed, light-resistant dispensing bottle. - To be administered with a metered-dose measuring device. |
| Auxiliary Labels | 1 Use as directed. Do not exceed prescribed dose. | 5 | Protect from light. |
| | 2 Keep out of reach of children. | 6 | Shake well before use. |
| | 3 Cap tightly after use. | 7 | Consult your health care practitioner if any prescription or over-the-counter medications are currently being used or are prescribed for future use. |
| | 4 Keep refrigerated. Do not freeze. | | |
| Pharmacist Instructions | Add any auxiliary labels specific to the API to the dispensing container as deemed necessary. | | |
| Patient Instructions | Contact your pharmacist in the event of adverse reactions. | | |



| | | | |
|-------------------|---|-----|-----------|
| Suggested Formula | Nystatin 100,000 IU/mL Oral Liquid (Suspension, 100 mL) | FIN | F 008 096 |
|-------------------|---|-----|-----------|

REFERENCES

| | |
|----|---|
| 1. | Glycerin. In: Rowe RC. <i>Handbook of Pharmaceutical Excipients, 4th Edition</i> . American Pharmaceutical Association; 2003: 257. |
| 2. | Xanthan Gum. In: Rowe RC. <i>Handbook of Pharmaceutical Excipients, 7th Edition</i> . American Pharmaceutical Association; 2012: 897. |
| 3. | Nystatin (Monograph). In: O'Neil MJ. <i>The Merck Index 14th Edition</i> . Whitehouse Station, NJ: Merck & Co, Inc.; 2006: Monograph #6736. |
| 4. | Nystatin (Monograph). <i>United States Pharmacopeia XLI / National Formulary 36</i> . Rockville, MD. US Pharmacopeial Convention, Inc. 2018: 2988. |
| 5. | Nystatin. In: Trissel LA. <i>Trissel's Stability of Compounded Formulations, 3rd Edition</i> . American Pharmaceutical Association; 2005: 317. |
| 6. | USP <795>. <i>United States Pharmacopeia XLI / National Formulary 36</i> . Rockville, MD. US Pharmacopeial Convention, Inc. 2018: 6546. |

DISCLAIMER: MEDISCA NETWORK INC., HEREBY REFERRED TO AS 'THE NETWORK', HAS PROVIDED THE FORMULA AND INSTRUCTIONS ABOVE AS A MODEL FOR EDUCATIONAL PURPOSES ONLY ON THE BASIS OF THE RECOGNIZED COMPENDIA AND TEXTS REFERENCED AT THE END OF THIS DOCUMENT. THE NETWORK TAKES NO RESPONSIBILITY FOR THE VALIDITY, SCHEDULING OR ACCURACY OF THIS INFORMATION OR FOR ITS SAFETY OR EFFECTIVENESS, NOR FOR ANY USE THEREOF, WHICH IS AT THE SOLE RISK OF THE LICENSED PHARMACIST. ADJUSTMENTS MAY BE NEEDED TO MEET SPECIFIC PATIENT NEEDS AND IN ACCORDANCE WITH A LICENSED PRESCRIBER'S PRESCRIPTION. THE PHARMACIST MUST EMPLOY APPROPRIATE TESTS TO DETERMINE THE STABILITY OF THIS SUGGESTED FORMULA. THE NETWORK CANNOT BE HELD LIABLE TO ANY PERSON OR ENTITY CONCERNING CLAIMS, LOSS, OR DAMAGE CAUSED BY, OR ALLEGED TO BE CAUSED BY, DIRECTLY OR INDIRECTLY, THE USE OR MISUSE OF THE INFORMATION CONTAINED IN THIS SUGGESTED FORMULA. IN ALL CASES IT IS THE RESPONSIBILITY OF THE LICENSED PHARMACIST TO KNOW THE LAW, TO COMPOUND ANY FINISHED PRODUCT AND TO DISPENSE THESE PRODUCTS IN ACCORDANCE WITH FEDERAL AND STATE LAW. MEDISCA NETWORK INC. MAKES NO WARRANTIES WITH RESPECT TO INFRINGEMENT OR NON-INFRINGEMENT BY THE FORMULA OF ANY PATENT OR OTHER INTELLECTUAL PROPERTY OF ANY OTHER PARTY, AND IT IS THE RESPONSIBILITY OF THE PHARMACIST TO INVESTIGATE AND DETERMINE ANY SUCH ISSUE.