



MEDISCA® NETWORK INC.
TECHNICAL SUPPORT SERVICES
FORMULATION CHEMISTRY DEPARTMENT
TOLL-FREE: 866-333-7811
TELEPHONE: 514-905-5096
FAX: 514-905-5097
technicalservices@medisca.net

4/14/2017; Page 1

| | | | |
|-------------------|---|-----|-------------|
| Suggested Formula | Bacitracin 400 U/mL, Neomycin 3.5 mg/mL, Polymyxin B Sulfate 5000 U/mL Topical Wound Liquid (Solution, 500 mL) | FIN | F 007 031v2 |
|-------------------|---|-----|-------------|

SUGGESTED FORMULATION

| Ingredient Listing | Qty. | Unit | NDC # | Supplier | Lot Number | Expiry Date |
|--|---------------|-------|-------|----------|------------|-------------|
| Bacitracin (Micronized), USP | 200 000 | Units | | | | |
| Neomycin Sulfate, USP | TBD | | | | | |
| Polymyxin B Sulfate, USP | 2 500 000 | Units | | | | |
| Sodium Chloride, USP | 3.97 | g | | | | |
| Benzalkonium Chloride Solution (50%), NF | 0.2 | mL | | | | |
| Sterile Water for Injection, USP | 400.0 | mL | | | | |
| Sterile Water for Injection, USP | q.s. to 500.0 | mL | | | | |
| Sodium Hydroxide 10% Solution | As required | | | | | |
| Hydrochloric Acid 10% Solution | As required | | | | | |



| | | | |
|-------------------|---|-----|-------------|
| Suggested Formula | Bacitracin 400 U/mL, Neomycin 3.5 mg/mL, Polymyxin B Sulfate 5000 U/mL Topical Wound Liquid (Solution, 500 mL) | FIN | F 007 031v2 |
|-------------------|---|-----|-------------|

SPECIAL PREPARATORY CONSIDERATIONS

Ingredient-Specific Information

| | |
|--|---|
| Light sensitive (protect from light whenever possible): | <i>Neomycin Sulfate, Polymyxin B Sulfate, Benzalkonium Chloride Solution</i> |
| Hygroscopic (protect from moisture whenever possible): | <i>Bacitracin, Neomycin Sulfate, Polymyxin B Sulfate Benzalkonium Chloride Solution</i> |
| Air sensitive (protect from air whenever possible): | <i>Benzalkonium Chloride Solution</i> |
| Metal reactive (do not allow to come into contact): | <i>Benzalkonium Chloride Solution</i> |
| Oxygen Sensitive (protect from oxygen whenever possible): | <i>Neomycin Sulfate</i> |

Suggested Preparatory Guidelines

☐ Non-Sterile Preparation ☒ Sterile Preparation

Processing Error / Testing Considerations: To account for processing error, pH testing sterility and endotoxin testing considerations during preparation, it is suggested to measure an additional **1 to 3%** of the required quantities of ingredients.

Special Instruction: This formula must be prepared within the appropriate facilities under adequate environmental conditions, following the necessary guidelines and procedures as stated within *USP 797*. Only trained and qualified personnel must prepare this formula.

All heat stable, reusable materials and equipment must be sterilized and depyrogenated by dry heat sterilization at 250°C for 2 hours prior to use.

Every batch of final product compounded using this procedure must be sterility and endotoxin tested before being dispensed.

Protective apparel, such as a sterile gown, sterile gloves, shoe covers, head cap, eyewear and face-masks should always be worn. In addition, proper personnel cleansing must be done before entering the buffer or clean area.

Filter integrity must be validated by performing a filter stress test. If the test demonstrates that the filter might be defective, the solution must be discarded and remade.

This procedure requires the use of very small quantities of ingredients. All calculations and preparation techniques must be verified before dispensing the final product.



| | | | |
|-------------------|---|-----|-------------|
| Suggested Formula | Bacitracin 400 U/mL, Neomycin 3.5 mg/mL, Polymyxin B Sulfate 5000 U/mL Topical Wound Liquid (Solution, 500 mL) | FIN | F 007 031v2 |
|-------------------|---|-----|-------------|

SUGGESTED PREPARATION (for 500 mL)

Weigh and / or measure the following ingredients when appropriate:

| Ingredient Listing | Qty. | Unit | Multiplication factor (*): ____ | Processing Error | Qty. to measure |
|--|---------------|-------|---------------------------------|------------------|-----------------|
| Bacitracin (Micronized), USP § | 200 000 | Units | | | |
| Neomycin Sulfate, USP § | TBD | | | | |
| Polymyxin B Sulfate, USP § | 2 500 000 | Units | | | |
| Sodium Chloride, USP § | 3.97 | g | | | |
| Benzalkonium Chloride Solution (50%), NF § | 0.2 | mL | | | |
| Sterile Water for Injection, USP § | 400.0 | mL | | | |
| Sterile Water for Injection, USP § | q.s. to 500.0 | mL | | | |
| Sodium Hydroxide 10% Solution § | As required | | | | |
| Hydrochloric Acid 10% Solution § | As required | | | | |

* Takes into account increased batch size conversions and density conversions, if required.

§ Weigh / measure just prior to use.

| Preparatory Instruction | |
|---|--|
| IMPORTANT: All preparatory procedures must be performed using proper Aseptic Technique | |
| 1. | <u>Equipment sterilization:</u> Following the manufacturer's specifications, sterilize and depyrogenate all heat stable, reusable materials and equipment, then return to ambient temperature. |



| | | | |
|-------------------|---|-----|-------------|
| Suggested Formula | Bacitracin 400 U/mL, Neomycin 3.5 mg/mL, Polymyxin B Sulfate 5000 U/mL Topical Wound Liquid (Solution, 500 mL) | FIN | F 007 031v2 |
|-------------------|---|-----|-------------|

2. **Ingredient quantification:**

A. Determine the potency of Bacitracin (Micronized) based on the certificate of analysis:

| | |
|--|-----------------------|
| | 100% |
| MINUS | |
| Loss on drying (from certificate of analysis) | _____ % |
| DIVIDED BY | 100 |
| EQUALS | |
| Quantity of dried Bacitracin (Micronized), in decimal | _____ |
| MULTIPLIED BY | |
| Assay on dried basis (from Certificate of Analysis) | _____ units/mg |
| EQUALS | |
| i. Potency of Bacitracin (Micronized) in (units/mg) | _____ units/mg |



| | | | |
|-------------------|---|-----|-------------|
| Suggested Formula | Bacitracin 400 U/mL, Neomycin 3.5 mg/mL, Polymyxin B Sulfate 5000 U/mL Topical Wound Liquid (Solution, 500 mL) | FIN | F 007 031v2 |
|-------------------|---|-----|-------------|

3. **Ingredient quantification:**

- A. Determine the quantity (in g) of Bacitracin required to make a 500 mL batch of Bacitracin 400 U/mL Topical Wound Liquid:

| | |
|--|--------------|
| Quantity of Bacitracin (Micronized) (in Units) required for 500 mL | 200 000 U |
| DIVIDED BY | |
| Bacitracin (Micronized) potency in (units/mg) (Step 2Ai) | _____ U/mg |
| EQUALS | |
| i Quantity of Bacitracin (Micronized) (in milligrams) needed for 500 mL | _____ mg |
| MULTIPLIED BY | |
| Multiplication factor – milligrams to grams | 0.001 |
| EQUALS | |
| ii Quantity of Bacitracin (Micronized) (in grams) needed for 500 mL | _____ g |
| MULTIPLIED BY | |
| Processing error adjustments (1 to 3%) | 1.01 to 1.03 |
| EQUALS | |
| iii Quantity of Bacitracin (Micronized) needed <i>plus</i> processing error adjustments | _____ g |



| | | | |
|-------------------|---|-----|-------------|
| Suggested Formula | Bacitracin 400 U/mL, Neomycin 3.5 mg/mL, Polymyxin B Sulfate 5000 U/mL Topical Wound Liquid (Solution, 500 mL) | FIN | F 007 031v2 |
|-------------------|---|-----|-------------|

4. **Ingredient quantification:**

A. Determine the potency of Neomycin Sulfate based on the certificate of analysis:

| | |
|--|-------------|
| MINUS | 100% |
| Loss on drying (from certificate of analysis) | _____ % |
| DIVIDED BY | 100 |
| EQUALS | |
| Quantity of dried Neomycin Sulfate, in decimal | _____ |
| MULTIPLIED BY | |
| Assay (base equivalent) on dried basis result (from certificate of analysis) | _____ µg/mg |
| MULTIPLIED BY (Multiplication factor – µg to grams /mg to grams) | 0.001 |
| EQUALS | |
| i. Potency of Neomycin Sulfate (Base equivalent) in g/g | _____ |



| | | | |
|-------------------|---|-----|-------------|
| Suggested Formula | Bacitracin 400 U/mL, Neomycin 3.5 mg/mL, Polymyxin B Sulfate 5000 U/mL Topical Wound Liquid (Solution, 500 mL) | FIN | F 007 031v2 |
|-------------------|---|-----|-------------|

5. **Ingredient quantification:**

- A. Determine the quantity (in g) of Neomycin Sulfate required to make a 500 mL batch of **Neomycin (Base)** 3.5 mg/mL Topical Wound Liquid:

| | |
|---|--------------|
| Quantity of Neomycin (Base) required for 500 mL | 1.750 g |
| DIVIDED BY | |
| Potency of Neomycin Sulfate (base equivalent), in g/g (Step 4Ai) | _____ |
| EQUALS | |
| i. Quantity of Neomycin Sulfate needed for 500 mL | _____ g |
| MULTIPLIED BY | |
| Processing error adjustments (1 to 3%) | 1.01 to 1.03 |
| EQUALS | |
| ii. Quantity of Neomycin Sulfate needed <i>plus</i> processing error adjustments | _____ g |



| | | | |
|-------------------|---|-----|-------------|
| Suggested Formula | Bacitracin 400 U/mL, Neomycin 3.5 mg/mL, Polymyxin B Sulfate 5000 U/mL Topical Wound Liquid (Solution, 500 mL) | FIN | F 007 031v2 |
|-------------------|---|-----|-------------|

6. **Ingredient quantification:**

A. Determine the potency of Polymyxin B Sulfate based on the certificate of analysis:

| | |
|--|-----------------------|
| | 100% |
| MINUS | |
| Loss on drying (from certificate of analysis) | _____ % |
| DIVIDED BY | 100 |
| EQUALS | |
| Quantity of dried Polymyxin B Sulfate, in decimal | _____ |
| MULTIPLIED BY | |
| Assay on dried basis (from Certificate of Analysis) | _____ units/mg |
| EQUALS | |
| i. Potency of Polymyxin B Sulfate in (units/mg) | _____ units/mg |



| | | | |
|-------------------|---|-----|-------------|
| Suggested Formula | Bacitracin 400 U/mL, Neomycin 3.5 mg/mL, Polymyxin B Sulfate 5000 U/mL Topical Wound Liquid (Solution, 500 mL) | FIN | F 007 031v2 |
|-------------------|---|-----|-------------|

7. **Ingredient quantification:**

- A. Determine the quantity (in g) of Polymyxin B Sulfate required to make a 500 mL batch of Polymyxin B Sulfate 5000 U/mL Topical Wound Liquid:

| | |
|--|--------------|
| Quantity of Polymyxin B Sulfate (in Units) required for 500 mL | 2 500 000 U |
| DIVIDED BY | |
| Polymyxin B Sulfate potency in (units/mg) (Step 6Ai) | _____ U/mg |
| EQUALS | |
| i Quantity of Polymyxin B Sulfate (in milligrams) needed for 500 mL | _____ mg |
| MULTIPLIED BY | |
| Multiplication factor – milligrams to grams | 0.001 |
| EQUALS | |
| ii Quantity of Polymyxin B Sulfate (in grams) needed for 500 mL | _____ g |
| MULTIPLIED BY | |
| Processing error adjustments (1 to 3%) | 1.01 to 1.03 |
| EQUALS | |
| iii Quantity of Polymyxin B Sulfate needed <i>plus</i> processing error adjustments | _____ g |



| | | | |
|-------------------|---|-----|-------------|
| Suggested Formula | Bacitracin 400 U/mL, Neomycin 3.5 mg/mL, Polymyxin B Sulfate 5000 U/mL Topical Wound Liquid (Solution, 500 mL) | FIN | F 007 031v2 |
|-------------------|---|-----|-------------|

| | |
|----|---|
| 8. | <p><u>Powder-liquid preparation:</u></p> <p>A. In the given order, sequentially add the following ingredients to the Sterile Water for Injection (400.0 mL <i>plus</i> processing error adjustments):</p> <ul style="list-style-type: none">-Sodium Chloride-Benzalkonium Chloride Solution (50%)-Bacitracin (amount determined in Step 3Aiii)-Neomycin Sulfate (amount determined in Step 5Aii)-Polymyxin B Sulfate (amount determined in Step 7Aiii) <p><u>Specifications:</u> Continuously mix until all solid particles have completely dissolved.</p> <p><u>End result:</u> Homogeneous liquid-like solution.</p> <p><u>Note:</u> Add the next ingredient, once the previous one has been completely added and dissolved.</p> |
| 9. | <p><u>pH testing:</u></p> <p>A. Draw an appropriate amount of the mixture (Step 8A).</p> <p>B. Test the pH of the sample. It should lie between 6.0 and 7.0.</p> <p>C. <u>If the pH < 6.0, carefully add, in a dropwise fashion, the Sodium Hydroxide 10% Solution to the mixture:</u></p> <ol style="list-style-type: none">1. Draw and transfer 1 or 2 drops of the Sodium Hydroxide 10% Solution to the mixture.2. Stir for at least 5 minutes to evenly disperse the Sodium Hydroxide 10% Solution.3. Re-test the pH.4. Continue to add the Sodium Hydroxide 10% solution until the pH of 6.0 to 7.0 is obtained. <p>IMPORTANT: Do not allow the pH to rise above 7.0.</p> <p>D. <u>If the pH > 7.0, carefully add, in a dropwise fashion, the Hydrochloric Acid 10% Solution to the mixture:</u></p> <ol style="list-style-type: none">1. Draw and transfer 1 or 2 drops of the Hydrochloric Acid 10% Solution to the mixture.2. Stir for at least 5 minutes to evenly disperse the Hydrochloric Acid 10% Solution.3. Re-test the pH.4. Continue to add the Hydrochloric Acid 10% Solution until the pH of 6.0 to 7.0 is obtained. <p>IMPORTANT: Do not allow the pH to fall below 6.0.</p> |



| | | | |
|-------------------|---|-----|-------------|
| Suggested Formula | Bacitracin 400 U/mL, Neomycin 3.5 mg/mL, Polymyxin B Sulfate 5000 U/mL Topical Wound Liquid (Solution, 500 mL) | FIN | F 007 031v2 |
| 10. | <u>Filling to volume:</u> A. Add additional Sterile Water for Injection to the above mixture to fill to the required batch size (500.0 mL <i>plus</i> processing error adjustments). <u>Specifications:</u> Continuously mix. <u>End result:</u> Homogeneous liquid-like solution. | | |
| 11. | <u>Filtering and transferring:</u> Aseptically filter the solution through a 0.22-µm sterile filter into the recommended dispensing container (see Packaging requirements). Transfer the remainder into a separate dispensing container. This is to be used as the Test sample for sterility and endotoxin testing. | | |
| 12. | <u>Filter integrity test:</u> Validate filter integrity by performing a filter stress test. If the test demonstrates that the filter might be defective, the solution must be discarded and remade. | | |
| 13. | <u>Sterility testing:</u> Validate the Test sample for sterility and endotoxins, in accordance to current USP 797 regulatory guidelines. | | |



| | | | |
|-------------------|---|-----|-------------|
| Suggested Formula | Bacitracin 400 U/mL, Neomycin 3.5 mg/mL, Polymyxin B Sulfate 5000 U/mL Topical Wound Liquid (Solution, 500 mL) | FIN | F 007 031v2 |
|-------------------|---|-----|-------------|

SUGGESTED PRESENTATION

| | | | | | |
|---------------------------|---|--|------------------------|--|--|
| Estimated Beyond-Use Date | | 7 days, refrigerated, as per USP 797. BUD based on a successful sterility and endotoxin test result. | Packaging Requirements | | Sterile, tightly closed, light-resistant dispersing bottle with sterile topical applicators. |
| Auxiliary Labels | 1 | Use as directed. Do not exceed prescribed dose. | 7 | Keep refrigerated. Do not freeze. | |
| | 2 | Keep out of reach of children. | 8 | Consult your health care practitioner if any prescription or over-the-counter medications are currently being used or are prescribed for future use. | |
| | 3 | Equilibrate to room temperature before use. | 9 | Cap tightly after use. | |
| | 4 | Do not use if product changes color. | 10 | Discard in the presence of particulate matter. | |
| | 5 | For topical use only. | 11 | Protect from light. | |
| | 6 | May impair mental and/or physical ability. Use care when operating a car or machinery. | 12 | Do not take with alcohol, sleep aids, tranquilizers or other CNS depressants. | |
| Pharmacist Instructions | | Add any auxiliary labels specific to the active ingredient to the dispensing container as deemed necessary. | | | |
| Patient Instructions | | Contact your pharmacist in the event of adverse reactions. IMPORTANT: The quantity of API administered is directly dependent on the quantity of product applied. | | | |



| | | | |
|-------------------|---|-----|-------------|
| Suggested Formula | Bacitracin 400 U/mL, Neomycin 3.5 mg/mL, Polymyxin B Sulfate 5000 U/mL Topical Wound Liquid (Solution, 500 mL) | FIN | F 007 031v2 |
|-------------------|---|-----|-------------|

REFERENCES

| | |
|-----|---|
| 1. | Parenteral Preparations. In: Allen, LV, Jr. <i>The Art, Science and Technology of Pharmaceutical Compounding Fourth Edition</i> . American Pharmaceutical Association; 2012: 341. |
| 2. | Benzalkonium Chloride. In: Rowe RC. <i>Handbook of Pharmaceutical Excipients, 7th Edition</i> . American Pharmaceutical Association; 2012: 60. |
| 3. | Sodium Chloride. In: Rowe RC. <i>Handbook of Pharmaceutical Excipients, 7th Edition</i> . American Pharmaceutical Association; 2012: 729. |
| 4. | Bacitracin. In: Sweetman SC, ed. <i>Martindale: The Complete Drug Reference, 36th Edition</i> . London, England: The Pharmaceutical Press; 2009: 210. |
| 5. | Neomycin Sulfate. In: Sweetman SC, ed. <i>Martindale: The Complete Drug Reference, 36th Edition</i> . London, England: The Pharmaceutical Press; 2009: 305. |
| 6. | Polymyxin B Sulfate. In: Sweetman SC, ed. <i>Martindale: The Complete Drug Reference, 36th Edition</i> . London, England: The Pharmaceutical Press; 2009: 317. |
| 7. | Bacitracin (Monograph). In: O'Neil MJ. <i>The Merck Index 15th Edition</i> . Whitehouse Station, NJ: Merck & Co, Inc.; 2013: Monograph #927. |
| 8. | Neomycin (Monograph). In: O'Neil MJ. <i>The Merck Index 15th Edition</i> . Whitehouse Station, NJ: Merck & Co, Inc.; 2013: Monograph #6539. |
| 9. | Polymyxin (Monograph). In: O'Neil MJ. <i>The Merck Index 15th Edition</i> . Whitehouse Station, NJ: Merck & Co, Inc.; 2013: Monograph #7693. |
| 10. | Bacitracin. In: Trissel LA. <i>Trissel's Stability of Compounded Formulations, 5th Edition</i> . American Pharmaceutical Association; 2012: 57. |
| 11. | Neomycin Sulfate. In: Trissel LA. <i>Trissel's Stability of Compounded Formulations, 5th Edition</i> . American Pharmaceutical Association; 2012: 349. |
| 12. | Polymyxin B Sulfate. In: Trissel LA. <i>Trissel's Stability of Compounded Formulations, 5th Edition</i> . American Pharmaceutical Association; 2012: 396. |
| 13. | Bacitracin (Monograph). <i>United States Pharmacopeia XXXIX / National Formulary 34</i> . Rockville, MD. US Pharmacopeial Convention, Inc. 2016: 2670. |
| 14. | Neomycin Sulfate (Monograph). <i>United States Pharmacopeia XXXIX / National Formulary 34</i> . Rockville, MD. US Pharmacopeial Convention, Inc. 2016: 5011. |
| 15. | Polymyxin B Sulfate (Monograph). <i>United States Pharmacopeia XXXIX / National Formulary 34</i> . Rockville, MD. US Pharmacopeial Convention, Inc. 2016: 5443. |
| 16. | USP <797>. <i>United States Pharmacopeia XXXIX / National Formulary 34</i> . Rockville, MD. US Pharmacopeial Convention, Inc. 2016: 626. |

DISCLAIMER: MEDISCA NETWORK INC., HEREBY REFERRED TO AS 'THE NETWORK', HAS PROVIDED THE FORMULA AND INSTRUCTIONS ABOVE AS A MODEL FOR EDUCATIONAL PURPOSES ONLY ON THE BASIS OF THE RECOGNIZED COMPENDIA AND TEXTS REFERENCED AT THE END OF THIS DOCUMENT. THE NETWORK TAKES NO RESPONSIBILITY FOR THE VALIDITY, SCHEDULING OR ACCURACY OF THIS INFORMATION OR FOR ITS SAFETY OR EFFECTIVENESS, NOR FOR ANY USE THEREOF, WHICH IS AT THE SOLE RISK OF THE LICENSED PHARMACIST. ADJUSTMENTS MAY BE NEEDED TO MEET SPECIFIC PATIENT NEEDS AND IN ACCORDANCE WITH A LICENSED PRESCRIBER'S PRESCRIPTION. THE PHARMACIST MUST EMPLOY APPROPRIATE TESTS TO DETERMINE THE STABILITY OF THIS SUGGESTED FORMULA. THE NETWORK CANNOT BE HELD LIABLE TO ANY PERSON OR ENTITY CONCERNING CLAIMS, LOSS, OR DAMAGE CAUSED BY, OR ALLEGED TO BE CAUSED BY, DIRECTLY OR INDIRECTLY, THE USE OR MISUSE OF THE INFORMATION CONTAINED IN THIS SUGGESTED FORMULA. IN ALL CASES IT IS THE RESPONSIBILITY OF THE LICENSED PHARMACIST TO KNOW THE LAW, TO COMPOUND ANY FINISHED PRODUCT AND TO DISPENSE THESE PRODUCTS IN ACCORDANCE WITH FEDERAL AND STATE LAW.